

Unhealthy food consumption among Ecuadorian children: A cross-sectional study in the context of the school food regulation

Juan Jácome¹ , Samuel Escandón² , Alejandro Rodríguez³ , Carl Lachat⁴ , Roberto Aguirre⁵ ,
Wilma Freire⁶ , René-Vinicio Sánchez⁷ , Silvana Donoso^{2,8} , Susana Andrade^{2,9} ,
Angélica Ochoa-Avilés^{2,8} .

Abstract: Unhealthy food consumption among Ecuadorian children: A cross-sectional study in the context of the school food regulation. **Introduction.** The intake of energy-dense unhealthy food at school could influence the overall energy intake of children. **Objective:** To characterize the patterns of buying a snack at school and to analyze the association of these patterns with the source of the school snack (brought from home or bought at school) and screen time. **Materials and methods:** A cross-sectional study was conducted from October 2018 until May 2019 among school children (9 to 12-year-olds) from Cuenca-Ecuador. Intake of energy-dense unhealthy food groups, the source of the school snack, and screen time were assessed using questionnaires. Multiple correspondence analysis was used to identify intake patterns of unhealthy food groups, and logistic regression to assess the association between intake of energy-dense unhealthy food groups (FG) with the school snack source. **Results:** Among 1,028 children recruited 63%, 42%, 30% and 22% of the children reported consuming sweets/confiture, savory snacks, fast food, and pastry products, respectively, at least two days per week. Around 39% of the children were classified as "Frequent consumers of energy-dense unhealthy FG". Buying the snack at school was associated with a more frequent intake of sweets/confiture (OR=1.56, CI 95% =1.05-2.32) and fast food (OR=2.01, CI 95% =1.15-3.50) during the week, as well as with being classified in the "frequent consumer of energy-dense unhealthy FG" (OR=1.99, 95% CI=1.40-2.82). **Conclusions:** Ecuadorian children still consume prohibited unhealthy foods at school. These results reinforce the importance of evaluating, monitoring, and adapting strategies to promote a balanced diet. **Arch Latinoam Nutr 2023; 73(3)S2: 35-46.**

Keywords: public policy, food quality, food legislation, school feeding, unhealthy food.

Resumen: Consumo de alimentos poco saludables en niños ecuatorianos: estudio transversal en el contexto del reglamento de alimentación escolar. **Introducción.** La ingesta de alimentos poco saludables con alto contenido energético en la escuela podría influir en la ingesta energética total de los niños. **Objetivo:** Caracterizar los patrones de compra de snacks en el colegio y analizar la asociación de estos patrones con la procedencia del snack escolar (de la casa o comprada en el colegio) y el tiempo de pantalla. **Materiales y métodos:** Se realizó un estudio transversal (octubre 2018-mayo 2019) en escolares (9-12 años) Cuenca- Ecuador. La ingesta de grupos de alimentos (FG) no saludables altos en energía, la fuente del snack escolar y el tiempo de pantalla se evaluaron mediante cuestionarios. Se utilizó análisis de correspondencias múltiples para identificar patrones de ingesta de FG poco saludables y la regresión logística para evaluar asociación entre la ingesta FG no saludables con la fuente del snack escolar. **Resultados:** En 1.028 niños 63%, 42%, 30% y 22% declararon consumir dulces/golosinas, aperitivos salados, comida rápida y productos de pastelería, respectivamente, al menos dos/días/ semana. Un 39% fueron clasificados como "Consumidores frecuentes de FG poco saludables de alta densidad energética". La compra de los snacks en el colegio se asoció con una ingesta más frecuente de dulces/golosinas (OR=1,56; IC95%=1,05-2,32) y comida rápida (OR=2,01; IC95%=1,15-3,50) durante la semana y con la clasificación en el grupo "consumidores frecuentes de FG poco saludables de alta densidad energética" (OR=1,99; IC95%=1,40-2,82). **Conclusiones:** Los niños ecuatorianos aún consumen alimentos no saludables prohibidos en la escuela. Estos resultados refuerzan la importancia de evaluar, monitorear y adaptar estrategias para promover una dieta balanceada. **Arch Latinoam Nutr 2023; 73(3)S2: 35-46.**

Palabras clave: política pública, calidad de alimentos, legislación alimentaria, alimentación escolar, comida no saludable.

¹Departamento de Ciencias de Alimentos y Biotecnología, Facultad de Ingeniería Química y Agroindustria, Escuela Politécnica Nacional, Quito 170525, Ecuador.

² Grupo Alimentación, Nutrición, Salud y Actividad Física, Departamento de Biociencias, Universidad de Cuenca. Cuenca, Ecuador. ³ Escuela de Medicina, Universidad Internacional del Ecuador, Quito, Ecuador. ⁴ Department of Food Technology, Safety and Health, Faculty of Bioscience Engineering, Universiteit Gent, Gent. ⁵ Carrera de Nutrición, Facultad de Ciencias Médicas, Universidad de Cuenca, Pje. del Paraíso, Cuenca, Ecuador. ⁶ Instituto de Investigaciones en Salud y Nutrición, Universidad San Francisco de Quito, Campus Cumbayá, Quito, Ecuador. ⁷ Grupo de Investigación y Desarrollo en Tecnologías Industriales (GIDTEC), Cuenca, Ecuador. ⁸ Carrera de Bioquímica y Farmacia, Universidad de Cuenca, Cuenca, Ecuador. ⁹ Carrera de Pedagogía de la Actividad Física y Deporte, Facultad de Filosofía, Universidad de Cuenca, Ecuador.

Autor para la correspondencia: Susana Andrade, e-mail: susana.andrade@ucuenca.edu.ec



Introduction

Obesity prevalence has increased worldwide (1); by 2016, around 340 million children and adolescents were overweight or obese (2). Unhealthy eating and sedentary behavior are important risk factors for obesity and non-communicable diseases (3). In Ecuador, 35% of school-age children are overweight or obese (4), 45% spend more than two hours/day on sedentary screen-related activities

(5), and risky dietary practices are prevalent in the adolescent population (6).

Healthy eating promotion requires broad and comprehensive policy actions (7). Food label regulations and food advertisement and marketing restrictions have been recognized as promising approaches (8,9). The last is especially important for children and adolescents, as screen-based sedentary behavior (i.e., TV viewing) is associated with consuming unhealthy energy-dense foods (10–12). In addition, setting standards and offering healthy foods in public institutions such as schools are promising strategies (13).

In Ecuador, two large-scale regulations are in place. Ecuador was the first Latin American country to implement food label standards in 2013. All processed and ultra-processed foods and beverages with added sugar, fat, or salt must carry a traffic-light-food label (14). The system classifies food items as low (green), medium (yellow), or high (red) in total fat, sugar, and salt content according to predefined concentrations (14). Secondly, the regulation of school food kiosks was issued in 2014. The kiosks must guarantee a variety of fresh and nutritious foods and beverages with a preference for traditional recipes; processed and ultra-processed foods and beverages with a red traffic-light, caffeinated, and energy drinks or drinks with non-caloric sweeteners are forbidden (15).

A priori impact evaluation of the regulations was not planned (15). The available evaluations have analyzed the implementation qualitatively or have quantified the (un)availability of (un)healthy foods in the school kiosks (16). Until now, no representative study has quantified which foods children buy at school kiosks; or has described the relationship between buying snacks at school and the intake of unhealthy energy-dense food groups.

The present study aims to: (i) characterize the overall intake of energy-dense unhealthy food groups (i.e., savory snacks, sweets, soft drinks, fast food, and pastry products) among schoolchildren; (ii) identify food groups that are usually consumed at school; and (iii)

evaluate if buying a snack at school is associated with the overall intake of unhealthy energy-dense food groups among schoolgoing children. The analysis will include important covariates (i.e., age, sex, weight status, socioeconomic level) and screen time as a predictor, considering that specific policies targeting food marketing and advertisement exposure among children are not in force.

Materials and methods

Design and context

A cross-sectional study involving children aged 9 to 12 was conducted in Cuenca from October 2018 until May 2019. The city is the third most populated in Ecuador (around 600,000 citizens), located in the Andean highlands (17). The Ecuadorian education system comprises public schools (the government fully subsidizes), partly subsidized schools (partly subsidized by the government and the parents), and private schools (parents cover the full tuition ranging from 70 to 400 USD/month) (18).

Sampling

This research is derived from a study that analyzed the individual and environmental factors associated with school children's health (19); therefore, the sample size of 934 was considered sufficient to detect these associations with a power of 85% and a cluster effect of two. A total of 20 schools were selected following a probability proportional to sample size procedure according to walkability index and neighborhood socioeconomic status of each school (20). In each selected school, 90 randomly students were invited to participate. All children with disabilities and pathologies that influence their usual diet or physical activity were excluded (flowchart in Figure 1).

Ethics statement

The Ethics Committee from Universidad San Francisco de Quito approved the survey instruments and the study protocol (No. 2017-090E, 2019-001E). All children and their parents/guardians signed written consent.

Data collection

Trained interviewers were in charge to apply questionnaires, following standardized procedures, that were adapted to the context and validated with children outside the study sample.




Figure 1. Flowchart of data collection

Socioeconomic level

Parents/guardians filled out the national survey on socioeconomic stratification (21) with a total score between 0 and 1000. A household was classified as low, medium and high socioeconomic status when the index was between 0-535, 536-696, and above 697, respectively.

Children's data

The child's ethnicity, age, sex, food intake, screen time, and anthropometric data was collected by using Kobo Toolbox (Harvard Humanitarian Initiative) on tablets.

Intake of energy-dense unhealthy food groups in the

previous week was assessed using a validated tool for Chilean children (22). Soft drink intake, including soda and processed and ultra-processed drinks, was reported as glasses consumed per day; the intake was categorized as <two glasses/week, ≥two glasses/day (23). Fast food (French fries, pizzas, hot dogs or hamburgers), savory snacks (cheese doodles, chips, nachos), sweets/confiture (ice creams, cookies, candies), and pastry products intake (cakes and sweet doughs) were reported as times of consumption per week and classified as: <two days/week, ≥two days/week (23). The source of the school snack was assessed using the following question:

Choose one option regarding the snack you eat at school during school hours: "You bring it from home, and you do not bring money", "You bring money to buy it", "You bring food from home and bring money to buy it", and "You do not eat a snack at school" (22). The daily snack allowance was registered for those who reported bringing money to buy snacks at school. The food groups usually consumed as school snacks were characterized for the children who reported "bringing their snack from home" and/or "buying it at school".

Screen time (min/day) after school hours during weekdays was measured using a Spanish version of the survey iHealth (24)(19). Total screen time in minutes included the time spent per day watching television/videos/DVD/Netflix, sedentary games (computer or consoles), and using the internet for leisure activities (social networks, YouTube). Screen time was categorized as inadequate when children spend > two hours and adequate for $\leq$ two hours (25,26).

Body weight and height were taken in duplicate (27) and a third measurement was taken in the case of a 2% difference. The arithmetic average of the two or three measurements was used to calculate children's body mass index (kg/m^2) and BMI z- (28,29).

Statistical analysis

Descriptive statistics are presented as percentages for categorical variables and mean $\pm$ standard deviation for continuous variables. Data management and analysis were performed using STATA (version 13, STATA Corp). All the analyses were adjusted for the cluster design using "svy" command. Hypothesis tests were performed with 0.05 alpha levels.

Food groups intake according to school snack source

According to its source (brought from home or bought at school), food groups consumed as school snacks (fruits, vegetables, savory snacks, sweets snacks, dairy, water, sugar drinks, non-sugar drinks, sandwiches, and

rice) were displayed using bar charts. Differences in food groups reported according to the source of the snack consumed at school were assessed using the test of two proportions.

Unhealthy food groups' intake patterns

Multiple correspondence analysis (MCA) was used to identify consumption patterns based on energy-dense unhealthy food groups (fast food, savory snacks, sweets/confiture, pastry products, and soft drinks). The technique analyses the relationships between several categorical dependent variables to produce a graphic illustration of the original information in a low dimensional space.

Five dimensions were explored, and two dimensions were chosen based on the alpha Cronbach coefficient, the percentage of variance explained, and associations between each variable with each dimension. A graphical display of food groups and subjects (children) was constructed using the two dimensions to identify consumption patterns. Having identified three consumption patterns of energy-dense unhealthy food groups intake, we used cluster analysis to classify children in each consumption pattern based on the z-values of MCA for the two selected dimensions. A scatter plot was constructed to outline how cluster analysis assigned children to each pattern. To confirm the MCA model's interpretability, cross tabulation frequencies and X2 test were used to identify clusters' differences in food group intake.

Predictors of daily or weekly intake of energy-dense unhealthy food groups

Logistic regression models were applied to assess if the school snack source and screen time were associated with the overall unhealthy food intake. Dependent variable in each model was the intake of sweets, savory snacks, fast food, pastry products, soft drinks, and the intake pattern of unhealthy food groups. For the latter variable, the three patterns of intake of unhealthy food groups (resulting in MCA analysis) were reduced to two groups ("Less frequent" and "Regular consumers" groups were merged into one category). The independent variables included school snacks source and screen time.

Also, the models were adjusted for age, sex, BMI z-scores, and socioeconomic level. Results are displayed as odds ratios (OR).

Results

A total of 1,028 children (48% attending private schools) were recruited. The average age of the children was 10.4 ± 1.2 years; 52% were female, and 75% identified themselves as mestizos. Also, 37%, 38%, and 25% of the children lived in high, medium, and low SES households, respectively. In 58% of the cases, the father was the head of the family, followed by the mother (35%). In total, 40% and 36% of the head of the family hold secondary and university or postgraduate degree, respectively. On average, the children spent 3.5 ± 2.3 hours/day on screen-time-related sedentary activities during school days, and 66.4% of children spent more than two hours on screen-time-related sedentary activities (Table 1).

Table 1. Participants characteristics (n=1028)

	n	%
Age (years); mean, SD	10.4	1.2
Female	533	51.9
Public school	527	51.26
Ethnicity		
Mestizos	766	74.51
White	119	11.58
Other ^a	143	13.91
Socio-economic status (n=833)		
Low	210	25.2
Medium	316	37.9
High	307	36.9
Head of the family (n=919)		
Father	534	58.1
Mother	320	34.8
Other ^b	65	7.1
Education level of the head of the family (n=915)		
Elementary	219	23.9
High school	365	39.9
University- postgraduate	331	36.2
Attend private school	501	48.7
BMI z-score; mean, SD	0.54	1.17
Screen time (Hours/day); mean, SD	3.5	2.3
Sedentary behavior (>2 hours)	683	66.44

^a Other ethnic minorities include indigenous, afro-descendant, montubio (mestizo from the Coastal region), mulatto (mixed black African with white European).

^b Other include grandfather, grandmother, uncle, aunt.

Energy-dense unhealthy food groups intake

Sweets/confiture and savory snacks intake for two or more days per week were reported by 63% and 42% of the children, respectively. In total, 30% and 22% of participants consumed fast food and pastry products at least twice weekly, respectively (Table 2).

The MCA and cluster analysis identified three groups (i.e., patterns) based on the intake of energydense unhealthy food groups (Figure 2a). The first pattern was named "Less frequent consumers" and was associated with those children who do not consume food from unhealthy food groups. The second

Table 2. Food groups intake and school snack source (n=1028)

Food groups intake and school snack source	n	%
Sweets and confiture		
<2 days/week	38.2	37.2
≥2 days/week	64.6	62.8
Savoury snacks		
<2 days/week	59.2	57.6
≥2 days/week	43.6	42.4
Fast food		
<2 days/week	71.5	69.6
≥2 days/week	31.3	30.4
Pastry products		
<2 days/week	80.0	77.8
≥2 days/week	22.8	22.2
Soft drinks		
<2 glasses/day	66.6	64.8
≥2 glasses/day	36.2	35.2
School snack source ^a		
Brought from home	17.4	16.9
Brought from home and bought at school	16.2	15.8
Bought at school	55.6	54.1
Not consume school snack	13.6	13.2
Snack allowance/day (USD); mean (SD), (n=718)	1.2	0.51

n=number of participants

^aSource of the food where children obtain their food




Figure 2. a) Multiple correspondence analysis with fast food intake, savory snacks intake, soft drinks intake, sweets and confiture intake, pastry products intake. b) Cluster analysis. No consumers n=208 (20.2%), Consumers group n=416 (40.5%), and Frequent consumers group n=404 (39.3%)

pattern was called "Regular consumers" and was associated with those children who consume fast food, savory snacks, sweets/confiture, and pastry products once a week and once a day for soft drinks. The last pattern was named "Frequent consumers" and was associated with those children who consume more than twice a week fast food, savory snacks, sweets/confiture, pastry products, and more than twice a day for soft drinks. Most of the children were clustered in the "Regular

consumers" (40.5%) and "Frequent consumers" (39.3%) groups, and 20.2% in the "Less frequent consumers" cluster (Figure 2b shows clusters according to MCA's Z values).

School snacks

In total 54% of children bought their snacks at school (n=718, average snack allowance was 1.2 ± 0.5 USD per day); 17% brought their snacks from home, and 16% combined both practices (i.e., both brought from home and bought at school) (Table 2).

The proportion of food groups consumed as school snacks by source (bought at school or brought from home) is presented in Figure 3. In summary, the proportion of children that reported having fruit, vegetable and dairy products as a snack were higher among those who brought from home the snack in comparison who bought at school (p -values <0.001). An important percentage of the participants reported the consumption of sweets and soft drinks as school snacks, although these percentage was higher among children who bought their snacks at school (sweets: 32% at school vs. 22% from home $p<0.001$; soft drinks 37% at school vs. 30% from home, $p=0.037$). There were no differences in the water ($p=0.349$) and savory snacks intake ($p=0.973$) between the children who

brought their snacks from home compared with those who bought them at school. Twenty-four percent of the children who reported bringing money to school indicated buying fast food (data unavailable for children bringing their snacks from home).

Associations between school snack source and screen time with a weekly or daily intake of energy-dense unhealthy food groups Buying the snack at school was associated with a more frequent intake of sweets/confiture (OR=1.56, 95% CI=1.05-2.32, $p<0.05$) and fast food (OR=2.01, 95% CI=1.15-3.50, $p<0.05$) over the week.




Figure 3. Proportion of intake of the food groups consumed as school snacks by source.
a) Nonprocessed food, b) Processed food; $n = 336$; home=brought from home, school=bought at school

Bringing money to buy snacks in the school kiosks was associated with being classified as the frequent consumer of energy-dense unhealthy food group (OR=1.99, 95% CI=1.40-2.82, $p<0.01$).

Spending two or more hours per day on screen devices was associated with a more frequent

intake of sweets (OR=2.13, 95% CI=1.54-2.94, $p<0.001$), savory snacks (OR=2.11, 95% CI=1.43-3.10, $p<0.01$) and fast food (OR=1.67, 95% CI=1.14-2.44, $p<0.05$). Finally, spending two or more hours per day on screens was independently associated with being classified as frequent consumer of energy-dense unhealthy food group (OR=1.67, 95% CI=1.18-2.36, $p<0.01$) (Table 3).

Table 3. Associations between sociodemographic characteristics, screen time and snack source

	High intake of sweets/confiture (≥ 2 days/week)	High intake of savory snacks intake (≥ 2 days/ week)	High intake of fast food (≥ 2 days/ week)	High intake of pastry products (≥ 2 days/week)	High intake of soft drinks (≥ 2 days/ week)	High intake of unhealthy food groups
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Snack source						
Brought from home (ref)	1.00	1.00	1.00	1.00	1.00	1.00
Brought from home and bought at school	1.48 (0.90-2.41)	0.80 (0.48-1.34)	1.70 (0.91-3.21)	1.56 (0.84-2.89)	1.07 (0.58-1.99)	1.30 (0.81-2.09)
Bought at school	1.56 (1.05-2.32) *	1.45 (0.94-2.22)	2.01 (1.15-3.50) *	1.50 (0.96-2.34)	1.11 (0.69-1.77)	1.99 (1.40-2.82) **
Screen time						
<2 hours/day (ref)	1.00	1.00	1.00	1.00	1.00	1.00
>2 hours/day	2.13 (1.54-2.94) ***	2.11 (1.43-3.10) **	1.67 (1.14-2.44) *	1.40 (0.88-2.24)	1.32 (0.94-1.85)	1.67 (1.18-2.36) **
BMI z-score	0.93 (0.84-1.03)	1.02 (0.91-1.15)	0.96 (0.81-1.13)	0.97 (0.78-1.21)	1.01 (0.88-1.16)	0.93 (0.82-1.05)
Age (years)	1.19 (1.01-1.41)*	1.06 (0.94-1.19)	1.20 (0.99-1.46)	0.97 (0.86-1.11)	1.03 (0.87-1.23)	1.10 (0.96-1.26)
Sex						
Girls (ref)	1.00	1.00	1.00	1.00	1.00	1.00
Boys	0.53 (0.39-0.73) **	1.17 (0.86-1.60)	1.13 (0.83-1.56)	0.76 (0.50-1.16)	1.15 (0.84-1.56)	0.89 (0.67-1.18)
Socioeconomic status						
Low (ref)	1.00	1.00	1.00	1.00	1.00	1.00
Medium	0.78 (0.48-1.27)	1.09 (0.66-1.79)	1.05 (0.63-1.74)	0.85 (0.48-1.53)	0.76 (0.52-1.09)	1.06 (0.65-1.73)
High	0.67 (0.40-1.13)	1.10 (0.83-1.45)	1.32 (0.83-2.12)	0.75 (0.42-1.34)	0.67 (0.43-1.04)	1.13 (0.76-1.68)

The model was fitted to BMI, age, sex and socioeconomic status. * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$

a Brought from home include children who reported bringing their snacks from home and do not bring money, and those who reported bringing food from home and also bringing money to buy it (n=336)

b Bought at school include children who reported bringing money to buy their snack exclusively (n=556)

Discussion

Our study documents how urban Ecuadorian children maintain unhealthy dietary behaviors that could be conducive to non-communicable diseases. Even though selling processed and ultra-processed foods and beverages with a red traffic light or with non-caloric sweeteners is forbidden at schools, our analysis suggests that buying food in the school kiosks is prevalent and associated with the intake of energy-dense unhealthy ultra-processed food groups over the week.

Two out of ten children reported buying fast food or savory snacks at school, three out of ten reported buying sweets and confiture and four out of ten bought soft drinks at school. The last can be an indirect indicator of the school food kiosks regulation compliance. The reasons for these results are diverse; according to the available data, the strategy seems not to be evaluated, updated or designed, considering the contextual and cultural aspects of the available evidence (15,30).

In this context, in April 2020, during the COVID-19 lockdown, the national parliament issued a national school food policy (30). The last demonstrated interest in the topic from policymakers.

Nevertheless, the policy seems not to have developed considering the local problems, barriers and facilitators, and local context in general. In addition, as far as we know, the policy was developed without the participation of local stakeholders and seemed not to be developed according to the available evidence of successful strategies. As a result, the strategies described are succinct. The policy focuses exclusively on food availability without considering other influential factors.

Furthermore, details about forbidden or allowed food items are not included; for example, nothing is mentioned about micronutrient content, type of fat, refined cereals, artificial sweeteners, processed and ultra-processed foods or food labels (30). Although in-person classes in the national school system started progressively in September 2021 (31), we could not find any additional update or plan to implement or evaluate the national food policy.

The large body of evidence must be considered to improve the strategies; in that sense, a recent systematic review of qualitative studies identified several barriers regarding the compliance of school food policies which seem not to be considered in the

current policy (32). The most common barriers include reductions in profit or revenues, availability of unhealthy food in the school proximities and, sociocultural aspects such as poor knowledge in the community or negative attitudes of local stakeholders towards the policies. These barriers have been previously identified in local qualitative and contextual research (16,33,34). Food kiosks owners often complain about the large availability of street vendors in the school surroundings or the poor preference of children to acquire fruit and vegetables, resulting in considerable food waste and profit losses. Besides, our data demonstrate that although the snacks sent from home seem to be healthier (16), a large proportion of parents (19-30%) sent savory snacks, sweets/confiture and soft drinks for the school snack. Previous studies performed in Cuenca demonstrated that poor parental knowledge, lack of time to prepare healthy snacks, children's financial autonomy, parental permissiveness, and unfavorable school support are influential factors for (un)healthy eating behaviors among schoolchildren and adolescents (35). None of these contextual and influential factors is considered in the school food policy.

Children in Ecuador are exposed to marketing and advertisement (6,36). Our results show that children spend an average of 3.5 h/day on sedentary screen activities, and 66.4% of children spend more than two hours on sedentary screen activities during school days. These figures seem to have increased, a study published in 2015 reported that Ecuadorian adolescents spent on average 2.2 h/day in sedentary activities (37), similarly, according to the national health survey performed in 2012, only 44.5% of children and adolescents exceeded two hours per day on screen-time sedentary activities (5).

Worryingly, our data suggest that spending more time on screen devices is associated with a pattern of intake of energy-dense ultra-processed unhealthy foods. Previous research indicated that sedentary screen time activities, such as high TV exposure, could be associated with a high intake of unhealthy food (38,39). The last could be explained due to non-conscious eating (40). Besides,

children are exposed to marketing during TV advertising, which promotes high-calorie products (10). These results highlight the urgent need to regulate food advertisements and marketing in TV, social media, and the school setting. According to the new school food policy, "The national health authority will coordinate with the competent Regulatory and Control Agency the restriction of advertising processed or ultra-processed foods and beverages with high salt, sugar and fat as well as those containing sweeteners". Once again, this is a good starting point that must be clarified to avoid ambiguity. For example, it is unclear if the traffic-light food label will be used to define the high content of salt, sugar and fat, and implementation details are not specified.

Undoubtedly, the school environment plays a crucial role in children's eating habits due to their time in school; for many children, at least two meals take place there (11). Our results highlight that purchasing food at school and sedentary screen time are potentially important and independent risk factors for unhealthy eating behaviors. Therefore, both factors should be the target of ongoing and future policy actions to enhance a more significant impact on children's health. One of the limitations of this study was that certain food groups were not considered (e.g., meat or fish) in the questionnaire used. Since this is a cross-sectional study, causation cannot be inferred. Nevertheless, considering the absence of formal evaluations, our results reasonably estimate the school food kiosks regulation compliance. We included a randomly selected representative sample of children living in urban areas; our data can be extrapolated to other urban settings in Ecuador..

Conclusions

This cross-sectional documents how Ecuadorian children maintain unhealthy food intake practices conducive to non-communicable diseases. Although the regulation of the school food kiosks had been in force, the school still seems to be a source

of intake of unhealthy food groups. Furthermore, screen time is strongly associated with unhealthy dietary patterns. More robust evidence-based policies are urgently needed to ensure that schools are healthy and supportive environments. Policymakers need to recognize the need to design culturally appropriate evidence-based strategies, which must be evaluated and adapted using good-quality research..

Acknowledgments

We are grateful to the parents, schools, students, authorities, and all the members of the project, especially to Elisabeth Rodas, Jorge Brito and Adriana Quezada for their insight and initial support. This work was developed within a joint postgraduate program of VLIR Network Ecuador.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Vazquez CE, Cubbin C. Socioeconomic Status and Childhood Obesity: a Review of Literature from the Past Decade to Inform Intervention Research. *Curr Obes Rep.* 2020; 9(4):562-570. <https://doi.org/10.1007/s13679-020-00400-2>
2. WHO. World Health Organization. 2021 [citado 16 de agosto de 2021]. Obesity and overweight. Disponible en: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
3. Guertin C, Pelletier L, Pope P. The validation of the Healthy and Unhealthy Eating Behavior Scale (HUEBS): Examining the interplay between stages of change and motivation and their association with healthy and unhealthy eating behaviors and physical health. *Appetite.* 2020;144:104487. <https://doi.org/10.1016/j.appet.2019.104487>
4. INEC. Salud, Salud Reproductiva y Nutrición [ENSANUT 2018 [Internet]. 2018 [citado 23 de marzo de 2021]. Disponible en: <https://www.ecuadorencifras.gob.ec/salud-salud-reproductiva-ynutricion/>
5. Andrade S, Ochoa-Avilés A, Freire W, Romero N, Orellana D, Contreras T, et al. Results From Ecuador's 2018 Report

- Card on Physical Activity for Children and Youth. *J Phys Act Health* 2018; 15 (S2): S344-S346. <https://doi.org/10.1123/jpah.2018-0536>
6. Ochoa-Avilés A, Verstraeten R, Lachat C, Andrade S, Van Camp J, Donoso S, et al. Dietary intake practices associated with cardiovascular risk in urban and rural Ecuadorian adolescents: A cross-sectional study. *BMC Public Health*. 2014; 14: 939. <https://doi.org/10.1186/1471-2458-14-939>
7. Hawkes C, Jewell J, Allen K. A food policy package for healthy diets and the prevention of obesity and diet-related non-communicable diseases: the NOURISHING framework. *Obes Rev*. 2013;14(S2):159-168. <https://doi.org/10.1111/obr.12098>
8. Story M, Nannery MS, Schwartz MB. Schools and obesity prevention: creating school environments and policies to promote healthy eating and physical activity. *Milbank Q*. marzo de 2009;87(1):71-100. <https://doi.org/10.1111%2Fj.1468-0009.2009.00548.x>
9. Wethington HR, Finnie RKC, Buchanan LR, et al. Healthier Food and Beverage Interventions in Schools: Four Community Guide Systematic Reviews. *Am J Prev Med*. 2020;59(1):e15-e26. <https://doi.org/10.1016/j.amepre.2020.01.011>
10. Webster EK, Staiano AE. Extended Heavy Television Viewing May Impact Weight Long Term in Adolescents. *J Adolesc Health*. 2020;66(5):517-519. <https://doi.org/10.1016/j.jadohealth.2020.02.007>
11. Clarke N, Pechey E, Mantzari E, et al. Impact of health warning labels on snack selection: An online experimental study. *Appetite*. 2020; 104744. <https://doi.org/10.1016/j.appet.2020.104744>
12. Lavriša Ž, Hristov H, Kelly B, Pravst I. Regulating children's exposure to food marketing on television: are the restrictions during children's programmes enough? *Appetite*. 2020;154:104752. <https://doi.org/10.1016/j.appet.2020.104752>
13. McHugh C, Hurst A, Bethel A, Lloyd J, Logan S, Wyatt K. The impact of the World Health Organization Health Promoting Schools framework approach on diet and physical activity behaviours of adolescents in secondary schools: a systematic review. *Public Health*. 2020;182:116-24. <https://doi.org/10.1016/j.puhe.2020.02.006>
14. Ministerio de Salud del Ecuador. Reglamento Sanitario De Etiquetado De Alimentos Procesados Para El Consumo Humano. Ecuador; 2013 p. 8.
15. Ministerios de Educación y Salud del Ecuador. Reglamento De Bases Escolares Del Sistema Nacional De Educación. Ecuador; 2014 p. 9.
16. Rivas-Mariño G. Análisis de la implementación de las normativas de oferta de alimentos y educación física en escuelas de Quito, 2019. Instituto Nacional de Salud Pública de México; 2021.
17. INEC. Encuesta Nacional de Empleo, Desempleo y Subempleo (ENEMDU), Diciembre 2019. Noticia. 2020;6.
18. Ministerio de Educación del Ecuador. Valores de pensiones y matrículas de las Instituciones Educativas regimen Sierra-Amazonia 2018-2019 [Internet]. 2018 [citado 10 de julio de 2020]. Disponible en: <https://educacion.gob.ec/wp-content/uploads/downloads/2018/08/ZONA6-SIERRA-AMAZONIA-2018-2019.pdf>
19. Molina-Cando MJ, Escandón S, van Dyck D, Cardon G, Salvo D, Fiebelkorn F, et al. Nature relatedness as a potential factor to promote physical activity and reduce sedentary behavior in Ecuadorian children. *PLoS ONE*. 2021;16(5): e0251972. <https://doi.org/10.1371/journal.pone.0251972>
20. IPEN. International Physical Activity and the Environment Network (IPEN) [Internet]. 2017 [citado 1 de junio de 2020]. Disponible en: http://ipenproject.org/methods_gis.html
21. INEC. 2011. 2011 [citado 8 de mayo de 2020]. p. 1-4 Encuesta de Estratificación del Nivel Socioeconómico NSE 2011. Disponible en: <https://www.ecuadorencifras.gob.ec/encuesta-deestratificacion-del-nivel-socioeconomico/>
22. Lera L, Fretes G, González CG, Salinas J, Vio F. Validación de un instrumento para evaluar consumo, hábitos y prácticas alimentarias en escolares de 8 a 11 años. *Nutr Hosp*. 2015;31(5):1977-1988. <https://dx.doi.org/10.3305/nh.2015.31.5.8607>
23. Pitt E, Cameron CM, Thornton L, et al. Dietary patterns of Australian children at three and five years of age and their changes over time: A latent class and latent transition analysis. *Appetite*. 2018; 129:207-216. <https://doi.org/10.1016/j.appet.2018.07.008>
24. Cerin E, Sit CHP, Huang YJ, Barnett A, Macfarlane DJ, Wong SSH. Repeatability of self-report measures of physical activity, sedentary and travel behaviour in Hong Kong adolescents for the iHealth(H) and IPEN – Adolescent studies. *BMC Pediatr*. 2014; 14:142. <https://doi.org/10.1186/1471-2431-14-142>
25. Dong Y, Lau PWC, Dong B, et al. Trends in physical fitness, growth, and nutritional status of Chinese children and adolescents: a retrospective analysis of 1.5 million students from six succes-

- sive national surveys between 1985 and 2014. *Lancet Child Adolesc Health*. 2019;3(12):871-880. [https://doi.org/10.1016/s2352-4642\(19\)30302-5](https://doi.org/10.1016/s2352-4642(19)30302-5)
26. Guthold R, Stevens GA, Riley LM, Bull FC. Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc Health*. 2020;4(1):23-35. [https://doi.org/10.1016/s2352-4642\(19\)30323-2](https://doi.org/10.1016/s2352-4642(19)30323-2)
27. Stewart A, Marfell-Jones M, Olds T, De Ridder J. *International Standards for Anthropometric Assessment*. ISAK 2011.
28. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ*. 2007;85(9):660-667. <https://doi.org/10.2471/blt.07.043497>
29. World Health Organization (WHO) 2019. BMI-for-age (5-19 years). Disponible en: https://www.who.int/growthref/who2007_bmi_for_age/en/
30. Asamblea Nacional Republica del Ecuador. *Ley Orgánica de Alimentación Escolar*. Ecuador; 2020 p. 24.
31. Ecuavisa. <https://www.ecuavisa.com/>. 2021 [citado 9 de noviembre de 2021]. En noviembre inicia el plan de retorno a clases presenciales de forma obligatoria. Disponible en: <https://www.ecuavisa.com/noticias/ecuador/en-noviembre-inicia-el-plan-de-retorno-a-clases-presenciales-de-forma-obligatoria-CD1027949>
32. Ronto R, Rath N, Worsley A, Sanders T, Lonsdale C, Wolfenden L. Enablers and barriers to implementation of and compliance with school-based healthy food and beverage policies: a systematic literature review and meta-synthesis. *Public Health Nutr*. 2020/04/22 ed. 2020;23(15):2840-2855. <https://doi.org/10.1017/s1368980019004865>
33. Freire WB, Waters WF, Rivas-Mariño G, Nguyen T, Rivas P. A qualitative study of consumer perceptions and use of traffic light food labelling in Ecuador. *Public Health Nutr*. 2016/09/13 ed. 2017;20(5):805-813. <https://doi.org/10.1017/s1368980016002457>
34. Angélica Ochoa GRM and RV. Tackling overweight and obesity in Ecuador: Policies and strategies for prevention. *Nutrition Exchange 8. Emergency Nutrition Network (ENN)*. 2017; 8: 12-14 <https://www.enonline.net/nex>
35. Verstraeten R, Van Royen K, Ochoa-Avilés A et al. A Conceptual Framework for Healthy Eating Behavior in Ecuadorian Adolescents: A Qualitative Study. *PLOS ONE*. 2014;9(1):e87183. <https://doi.org/10.1371/journal.pone.0087183>
36. Verstraeten R, Leroy J, Pieniak Z, Ochoa-Avilés A, Holdsworth M, Verbeke W, et al. Individual and Environmental Factors Influencing Adolescents' Dietary Behavior in Low- and Middle- Income Settings. *PLOS ONE*. 2016; 11: e0157744. <https://doi.org/10.1371/journal.pone.0157744>
37. Andrade S, Verloigne M, Cardon G. et al. School-based intervention on healthy behaviour among Ecuadorian adolescents: effect of a cluster-randomized controlled trial on screen-time. *BMC Public Health*. 22 de septiembre de 2015;15:942. <https://doi.org/10.1186/s12889-015-2274-4>
38. Higgs S. Manipulations of attention during eating and their effects on later snack intake. *Appetite*. 2015;92: 287-294. <https://doi.org/10.1016/j.appet.2015.05.033>
39. Shqair AQ, Pauli LA, Costa VPP, Cenci M, Goettems ML. Screen time, dietary patterns and intake of potentially cariogenic food in children: A systematic review. *J Dent*. 2019;86:17-26. <https://doi.org/10.1016/j.jdent.2019.06.004>
40. Tapper K, Seguias L. The effects of mindful eating on food consumption over a half-day period. *Appetite*. 2020;145:104495. <https://doi.org/10.1016/j.appet.2019.104495>