

Insulin-like growth factor-1 promotes the proliferation and odontogenic differentiation of human dental pulp cells *in vitro* and *in vivo*.

Yan Wang, Nan Du, Cong-na Liu and Wen-jing Li

Department of Oral Medicine, The Second Hospital of Hebei Medical University, Shijiazhuang, China.

Keywords: dental tissue regeneration; odontoblasts; osteogenic differentiation.

Abstract. Human dental pulp cells (hDPCs) have emerged as a potential alternative for the regeneration of dental tissues. Insulin-like growth factor-1 (IGF-1) is involved in the proliferation and osteogenic differentiation of hDPCs *in vitro*. However, the effect of IGF-1 on the proliferation and odontogenic differentiation of hDPCs *in vivo* remains unknown. This study collected hDPCs from healthy premolars and third molars by collagenase type I and dispase. Immunocytochemical staining showed positive vimentin staining and negative cytokeratin staining in hDPCs. Treatment with IGF-1 (50, 75, and 100 ng/mL) significantly increased the proliferation ability of hDPCs in a concentration-dependent manner. *In vivo* experiments, hDPCs were seeded into an acellular dermal matrix and transplanted subcutaneously into nude mice. After two and four weeks of transplantation, the hematoxylin and eosin staining revealed more cells and extracellular matrix in implants from the IGF-1 treatment group, and Alizarin Red staining revealed more mineralized tissue compared to the control group. Transmission electron microscopy (TEM) analysis of hDPCs showed an abundance of mitochondria, rough endoplasmic reticulum, and Golgi complexes. In conclusion, IGF-1 promotes the proliferation of hDPCs *in vitro* and odontogenic differentiation of hDPCs *in vivo*, indicating that modifying IGF-1 signaling may provide potential strategies for the regeneration of dental tissues.

El factor de crecimiento similar a la insulina-1 promueve la proliferación y diferenciación odontogénica de células de pulpa dental humana *in vitro* e *in vivo*.

Invest Clin 2024; 65 (2): 179 – 191

Palabras clave: regeneración del tejido dental; odontoblastos; diferenciación osteogénica.

Resumen. Las células de la pulpa dental humana (hDPCs) han emergido como una alternativa prometedora para la regeneración de tejidos dentales. El factor de crecimiento insulínico tipo 1 (IGF-1) juega un rol crucial en la proliferación y diferenciación osteogénica de las hDPCs en condiciones *in vitro*. No obstante, el impacto del IGF-1 sobre la proliferación y diferenciación odontogénica de las hDPCs en un contexto *in vivo* aún no ha sido completamente elucidado. En el presente estudio, se extrajeron hDPCs de premolares y terceros molares sanos mediante el uso de colagenasa tipo I y dispasa. La tinción inmunocitoquímica de las hDPCs reveló una reactividad positiva para vimentina y negativa para citoqueratina. El tratamiento con IGF-1 en concentraciones de 50, 75 y 100 ng/mL incrementó de manera significativa y dependiente de la dosis la capacidad proliferativa de las hDPCs. En experimentos *in vivo*, las hDPCs fueron implantadas en una matriz dérmica acelular y posteriormente trasplantadas de manera subcutánea en ratones desnudos. Tras 2 y 4 semanas de trasplante, la coloración con hematoxilina y eosina evidenció un aumento en la cantidad de células y matriz extracelular en los implantes tratados con IGF-1, mientras que la coloración con Rojo de Alizarina indicó una mayor formación de tejido mineralizado en comparación con el grupo control. El análisis mediante microscopía electrónica de transmisión (TEM) de las hDPCs mostró una abundante presencia de mitocondrias, retículo endoplásmico rugoso y complejos de Golgi. En conclusión, nuestros hallazgos sugieren que el IGF-1 favorece la proliferación de las hDPCs *in vitro* así como su diferenciación odontogénica *in vivo*, lo cual señala que la modificación de la señalización del IGF-1 podría ofrecer estrategias potenciales para la regeneración de tejidos dentales.

Received: 24-10-2023 *Accepted:* 02-03-2024

INTRODUCTION

Dental pulp, situated at the innermost tissue within teeth, primarily comprises loose connective tissue and plays an essential role in the repair and regeneration of dental tissues. Traumatic incidents or the impact of infectious agents and other pathogenic stimuli can readily give rise to pulpitis¹. Human dental pulp cells (hDPCs) are

present within dental pulp tissue, constituting a mixture of fibroblasts, inflammatory, immune cells, odontoblasts, and undifferentiated mesenchymal cells^{2,3}. Among these components, the undifferentiated mesenchymal cells possess a multipotent cellular phenotype with the potential to differentiate into various cell lineages, such as osteoblasts, adipocytes, neural progenitors, and chondrocytes⁴⁻⁶. When dental pulp tissue is

exposed to external stimuli, dental pulp cells can differentiate into odontoblasts under the regulation of various factors, which is a significant factor in the regeneration and repair capacity of dental tissue damage^{7,8}. Therefore, exploring the mechanisms of hDPCs proliferation and odontoblastic differentiation is essential.

Cell-based tissue engineering has become an irreplaceable method for regenerating dental tissues⁹. Its fundamental mechanism involves inducing stem cells to differentiate into odontoblast-like or osteoblast-like cells, ultimately achieving the regeneration of the dental pulp-dentin complex^{10,11}. Furthermore, hDPCs exhibit some advantages, characterized by no ethical controversy, low immunogenicity, and easy obtainment from impacted or orthodontic extraction^{12,13}. Therefore, hDPCs have emerged as a potential alternative for future use in regeneration therapy. In addition to stem cells and biomimetic materials, suitable growth factors are imperative in tissue regeneration by promoting cell proliferation, differentiation, and locomotion^{14,15}.

Insulin-like growth factor-1 (IGF-1), a member of the insulin-like peptide family, exerts essential roles in the bone formation and periodontal regeneration of teeth^{14,16,17}. Local controlled delivery of IGF-I from dextran-co-gelatin hydrogel microspheres enhances new bone formation and alveolar bone reconstruction in the periodontal defects¹⁷. Moreover, IGF-1 is also involved in the odontoblastic differentiation of hDPCs or human dental pulp stem cells (hDPSCs). In vitro experiments showed 100 ng/mL IGF-1 promotes proliferation and increases the osteogenic differentiation-related expression of DPSCs via mammalian target of rapamycin (mTOR) signaling pathway¹⁸, as well as the enhancement of proliferation and osteo/odontogenic differentiation was found in the human periodontal ligament stem cells via activating MAPK pathways¹⁹. Similarly, other in vitro research indicates that combined use of 100 ng/ml VEGF and 100 ng/mL IGF-

1 also improve the proliferation, migration, osteogenesis, and angiogenesis of hDPSCs via activation of the phosphoinositide 3-kinase (PI3K)/Akt signaling pathway²⁰.

This study aimed to investigate the effect of IGF-1 on the proliferation and odontogenic differentiation of hDPCs *in vitro* and *in vivo*. These findings may offer valuable insights into the potential utility of hDPCs in treating dental pulp diseases.

MATERIALS AND METHODS

Isolation of human DPCs

Healthy premolars for orthodontic needs or third molars were collected from patients without dental caries or periodontal tissue diseases at the dental clinic of The Second Hospital of Hebei Medical University. All study protocols were approved by the Ethics Committee of The Second Hospital of Hebei Medical University, and written informed consent was obtained from all patients. The hDPCs were obtained as described previously. To completely collect the dental pulp, the tooth surface attachments were scraped off, and the teeth were split in a sterile environment. After washing three times with phosphate buffer solution (PBS), dental pulp tissues were cut into pieces of 1 mm×1 mm in size, then digested in a solution of 3mg/mL collagenase type I (Sigma-Aldrich) and 4mg/mL dispase (Sigma-Aldrich) at 37°C on a shaker for one hour. Following filtration and centrifuging, cell suspensions of dental pulp were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco, Grand Island, NY, USA) supplemented with 20% fetal bovine serum and 1% penicillin-streptomycin (Gibco, Grand Island, NY, USA) at 37°C in a 5% CO₂ incubator. Observe whether dental pulp cells crawl out of the tissue blocks and cell morphology and growth. Once cells adhere to the culture surface, the medium was refreshed every three days. When cell fusion reached over 80%, the adherent cells were digested with 0.25% trypsin (Sigma-Aldrich), and passage culture was performed at a 1:3

ratio. The hDPCs from the mixed population at passages 3 to 6 were used for subsequent experiments.

Immunohistochemical staining

Passage 4 hDPCs were digested with 0.25% trypsin and prepared as cell suspension, then seeded into 6-well cell culture plates at a concentration of 1×10^4 cells per well. Immunohistochemical staining for vimentin and cytokeratin was carried out on the day following cell adherence. Briefly, cell slides were fixed with 4% formaldehyde for 10 minutes, then treated with 0.5% Triton X-100 (diluted in PBS) for 10 minutes at room temperature. Following PBS washing, cells were blocked with 10% FBS for 20 minutes and subsequently incubated overnight at 4°C with primary antibodies anti-vimentin and anti-cytokeratin (1:200, Abcam). After washing three times with PBS, cells were incubated with biotin-conjugated secondary antibodies (1:1000, Abcam) at room temperature for 20 minutes. DAB staining was performed, and counterstaining with hematoxylin was carried out to visualize the cell nuclei. After mounting coverslips with neutral balsam, photography was conducted under an optical microscope (Olympus, Tokyo, Japan).

Flow cytometry analysis

The purity of isolated hDPCs was assessed using flow cytometry analysis with specific antibodies for CD44 and CD45²¹. The cells were trypsinized and fixed in 4% paraformaldehyde in PBS. Subsequently, hDPCs were permeabilized with 0.1% Triton X-100 (Sigma-Aldrich) in PBS and then incubated with conjugated antibodies against CD44 (1:100, Biotech, Minneapolis, USA) and anti-CD45 (1:100, Biotech, Minneapolis, USA). After washing three times with PBS, flow cytometric analysis was conducted using a FACS (Becton, Dickinson, San Jose, CA) with Quest CELL software (Becton, Dickinson).

Cell proliferation assay

The proliferation ability of hDPCs was assessed using the MTT (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl nyltetrazolium bromide, Sigma) assays. Passage 4 hDPCs were prepared as a cell suspension with 0.25% trypsin and seeded into 96-well cell culture plates at a concentration of 2×10^3 cells per well. After 24 hours of cultivation, cells were randomly divided into four groups. In the experimental groups, DMEM culture medium containing different concentrations of IGF-1 (50 ng/mL, 75 ng/mL, 100 ng/mL, Peprotech, USA) was added, while in the control group, DMEM without IGF-1 was added. Each group had five replicates. The cells were then cultured at 37°C for 1, 3, 5, and 7 days, respectively. Subsequently, 20 μ L of MTT (2 mg/mL in PBS) was added to each well. After 4 hours of incubation, the supernatant was removed, and 150 μ L DMSO were added. The absorbance values at A450 for each well were determined using a microplate reader (Bio-Rad).

Odontoblastic differentiation *in vivo*

The acellular dermal matrix (ADM, Beijing Qingyuan Weiye Company, China) underwent aseptic cutting into pieces measuring 5mm \times 5mm \times 1mm. These pieces were subjected to three washings with PBS and a 2-hour soak in 10% FBS within a sterile environment. The 2×10^5 hDPCs were prepared as a cell suspension. After removing the FBS, cells were gently seeded onto the scaffold using a pipette, ensuring even distribution within the scaffold interstices through gentle shaking. The composite was co-cultured at 37°C for 24 hours. The culture medium was replaced with 100 ng/mL IGF-1 culture medium supplemented with mineralization induction solution, consisting of 10% FBS, DMEM, 10 mmol/L β -glycerophosphate, 100 mg/mL vitamin C, and 10 nmol/L dexamethasone. The culture medium in the control group was replaced with a mineralization induction solution without IGF-1. The

induction culture was maintained for three days. For the *in vivo* phase, eight 4-week-old immunodeficient mice were selected. Under sterile conditions, incisions were made in the skin along their dorsal midlines. The hDPCs-scaffold complexes were transplanted subcutaneously into the left dorsal region, while control scaffolds were transplanted as controls into the right dorsal region. The odontoblastic differentiation in the hDPCs-scaffold complexes within the mice was observed at two and four weeks post-transplantation.

Histological staining

Hematoxylin and Eosin (HE) staining was used for the histological observation, and Alizarin red staining was performed to identify calcium-containing osteocytes or odontoblasts. The mice were euthanized at two and four weeks post-transplantation. The scaffold and surrounding tissues were immediately harvested and fixed in 4% paraformaldehyde. Subsequently, 4- μ m-thick sections were prepared using a freezing microtome. The sections were stained with hematoxylin for 3 minutes for HE staining, followed by differentiation and a 20-second rinse in alkaline water. Next, sections were stained with eosin for 20 seconds. For the alizarin red staining, sections were fixed in 95% anhydrous ethanol for 10 minutes. After rinsing in double-distilled water thrice, sections were stained in 0.1% alizarin red-Tris-HCl at 37°C for 30 minutes. All sections were observed and counted under a light microscope (Olympus, Tokyo, Japan).

Transmission electron microscopy (TEM)

TEM was used to assess the morphologies of hDPCs-scaffold complexes. After two and four weeks post-transplantation, the scaffold was immediately harvested and washed three times with PBS, then fixed in 4% (v/v) glutaraldehyde for 2 hours at room temperature. The complexes were dehydrated with varying ethanol concentrations and embedded with epoxy resin. Subse-

quently, 4- μ m-thick sections were prepared and stained with the toluidine blue staining. Ultimately, all the samples were air-dried overnight and viewed with a Hitachi Model H-7500 TEM (Hitachi, Japan).

Statistical analysis

Data were shown as mean \pm SD, and statistical analysis was performed using SPSS 17.0 software (IBM SPSS, Armonk, NY, USA). Group comparisons were performed using one-way ANOVA followed by a least significant difference (LSD) post-hoc comparisons. $P < 0.05$ was considered to indicate a statistically significant difference.

RESULTS

Identification of human dental pulp cells (hDPCs)

The inverted microscopy showed that hDPCs displayed clustered growth after seven days of adherent culture, primarily with a spindle-shaped morphology, while a smaller fraction of polygonal shapes (Fig. 1A). After passaging, the hDPCs adopted a vortex-like growth pattern, characterized by a more uniform, short spindle-shaped morphology with extending cytoplasmic processes (Fig. 1B). Immunocytochemical analysis was performed using passage 4 hDPCs, and showed positive staining for vimentin (Fig. 1C) and negative staining for cytokeratin (Fig. 1D), indicating their mesenchymal origin. Additionally, the flow cytometry analysis showed that the purity of hDPCs reached more than 95% (Fig. 1E). These morphological features were aligned with the typical biological characteristics of hDPCs.

IGF-1 promoted proliferation of hDPCs *in vitro*

MTT assay was performed to assess the effect of different IGF-1 concentrations (50, 75, and 100 ng/mL) on the proliferation ability of hDPCs. As shown in Fig. 2, there was no significant difference among groups ($p > 0.05$), and 100 ng/mL IGF-1 sig-

nificantly increased the proliferation of hDPCs three days after treatment compared to the control group ($p < 0.01$). At day five and day seven, treatment with IGF-1 significantly increased the proliferation of hDPCs compared with the control group ($p < 0.05$

and $p < 0.01$), 100 ng/mL IGF-1 displayed a more substantial enhancement compared to 50 ng/mL and 75 ng/mL ($p < 0.05$ and $p < 0.01$). These results indicated that IGF-1 promoted the proliferation of hDPCs in a concentration-dependent manner.




Fig. 1. Identification of hDPCs. A: Morphology after seven days of primary culture. B: Morphology of hDPCs in passage 3. C: Immunocytochemical staining for vimentin. D: Immunocytochemical staining for cytokeratin. E: Flow cytometric analysis for CD44 and CD45.




Fig. 2. IGF-1 promoted the proliferation of hDPCs in vitro. MMT assay showed treatment with 100 ng/mL IGF-1 significantly increased the proliferation of hDPCs from day 3, as well as 50 ng/mL and 75 ng/mL IGF-1 in day 5 and day 7 cultures. * $p < 0.01$ and ** $p < 0.01$ vs control (DMEM without IGF-1).

IGF-1 promoted odontogenic differentiation of hDPCs *in vivo*

In order to investigate the effect of IGF-1 on osteogenic differentiation *in vivo*, hDPCs were seeded into an ADM scaffold for the induction of odontogenic differentiation, and the hDPCs-scaffold complexes were transplanted subcutaneously into immunodeficient mice for two or four weeks. HE staining conducted at 2 and 4 weeks post-transplantation revealed more cells and extracellular matrix in implants from IGF-1 treatment group compared to the control group (Fig. 3). Alizarin red staining demonstrated that mineralized nodules in the IGF-1 group exhibited time-dependent enhancement, including the differentiation of odontoblast-like cells (Fig. 4). TEM analysis of hDPCs at 2 and 4 weeks post-transplantation showed an abundance of mitochondria, rough endoplasmic reticulum and Golgi complexes (Fig. 5). These findings indicated that IGF-1 progressively promoted the odontogenic differentiation of hDPCs *in vivo*.

DISCUSSION

Cell-based tissue engineering has become an irreplaceable method for regenerating dental tissues⁹, and the potential value of hDPCs has been widely accepted in bone tissue engineering due to their high self-renewal capacity and stemness^{22,23}. The process of dentinal regeneration involves the proliferation and differentiation of hDPCs into odontoblasts, dental pulp healing, and reparative dentin formation²⁴. Increasing studies have implicated IGF-1 in the maintenance of proliferation and differentiation of various stem cells, such as embryonic stem cells, bone marrow mesenchymal stem cells (BMSC), periodontal ligament stem cells, and hDPSCs²⁵. However, the effect of IGF-1 on the proliferation and differentiation of hDPCs *in vitro* and *in vivo* warrants further investigation. In this study, we found that exogenous IGF-1 promoted the proliferation of hDPCs in a concentration-dependent manner.




Fig. 3. HE staining revealed more cells and extracellular matrix in implants from the IGF-1 group compared to the control group (without IGF-1) at 2 and 4 weeks post-transplantation.




Fig. 4. IGF-1 promoted odontogenic differentiation of hDPCs *in vivo*. Alizarin red staining was performed to identify calcium-containing odontoblast, showing more mineralized tissues (arrows) in IGF-1 group compared to the control group (without IGF-1) at 2 and 4 weeks post-transplantation.




Fig 5. TEM analysis. A-B: The subcellular structure of hDPCs in scaffold 2 weeks (A) and 4 weeks (B) post-transplantation in IGF-1 group (magnification $\times 6K$). C-D: The subcellular structure of hDPCs in scaffold 2 weeks (C) and 4 weeks (D) post-transplantation in IGF-1 group (magnification $\times 15K$). Mitochondria and rough endoplasmic reticulum were highlighted using arrows.

Additionally, IGF-1 progressively promoted odontogenic differentiation of hDPCs *in vivo* by subcutaneous transplantation into nude mice.

As a unique type of dental stem cell, hDPCs are derived from the pulp in deep carious teeth and display stronger proliferation and osteogenic differentiation capability due to distinctive environmental stimulus^{26,27}. This distinctive trait provides valuable insights into the cellular mechanisms underlying tooth development and regenerative repair, offering the potential for dental pulp regeneration. The proliferation ability of colony-forming cultures of hDPCs and BMSCs was assessed by bromodeoxyuridine, showing that the number of proliferating cells in hDPC cultures was significantly higher than that in BMSC cultures. This result proves the more substantial proliferation capability of hDPCs compared to BMSCs²⁸. Moreover, under specific induction conditions, hDPCs can be induced to differentiate into various cell types, including odontoblast-like and osteoblast-like cells. Almushayt *et al.*²⁹ experimentally verified the capacity of hDPCs to differentiate into odontoblast-like cells *in vitro*. This study also found the odontoblast differentiation potential of hDPCs *in vivo* by subcutaneous transplantation into nude mice. Zhang *et al.*³⁰ isolated dental pulp cells from healthy third molars through enzymatic digestion, confirming their identity as hDPSCs. Subsequent induction assays revealed their multilineage differentiation potential, including differentiation potential toward adipogenic, osteogenic, fibrogenic, chondrogenic, and neurogenic lineages. Yang *et al.*³¹ demonstrated the capacity of dental pulp stem cells to differentiate into chondrocytes and myocytes under specific induction conditions.

Growth factors are critical in tooth tissue regeneration, as they have essential roles in regulating cell functions, such as platelet-derived growth factor, fibroblast growth factor, and epidermal growth factor^{32,33}. IGF-1, one of the ubiquitous peptide hormones, has

been identified to facilitate various cellular processes, including cell proliferation, differentiation, migration, apoptosis, and survival²⁵. Previous studies have reported that IGF-1 can affect hDPCs proliferation, promote osteogenesis, and help reconstruct tooth-supporting tissues³⁴⁻³⁶. A report by Onishi *et al.*³⁷ reveals multiple biological effects of IGF-1 on hDPCs. IGF-1 can stimulate hDPCs proliferation, increase mucin and extracellular matrix protein synthesis, enhance DNA synthesis, and increase alkaline phosphatase (ALPase) activity when hDPCs were cultured in a serum-free medium supplemented with IGF-1. Increasing investigations have illustrated the capacity of IGF-1 to induce the differentiation of ameloblasts and odontoblasts, facilitating dentin regeneration and the development of structures resembling enamel-dentin complexes^{34,35}. IGF-1 also influences the proliferation and differentiation of other odontogenic cell types, specifically periodontal ligament stem cells and stem cells from apical papilla^{19,38}. In this study, we also found that exogenous IGF-1 promoted the proliferation of hDPCs *in vivo* in a concentration-dependent manner and progressively promoted odontogenic differentiation of hDPCs *in vivo*.

IGF-1 receptor (IGF-1R) is a cell-surface receptor tyrosine kinase with its specific ligands IGF-1³⁹. It activates downstream signaling pathways, namely the phosphoinositide 3-kinase (PI3K)/AKT and the RAS/mitogen-activated protein kinase (MAPK) pathways. These pathways are widely involved in stem cell growth, proliferation, and differentiation^{40,41}. Previous studies have reported that the activation of IGF-1R signaling contributes to maintaining the self-renewal and differentiation of hDPCs²⁵. It has been demonstrated that treatment with IGF-1 can up-regulate the expression of phospho-ERK and phospho-p38 of hDPCs, suggesting MAPK signaling pathway during differentiation of hDPSCs and human periodontal ligament stem cells¹⁹. The interactions between IGF-1R and the p38 MAPK signaling pathway

control the quiescence and activation of hDPSCs. In addition, the downstream effectors between the MAPK signaling pathway and the PI3K/Akt signaling pathway partly overlap, resulting in the interaction effect of two signaling pathways. Some studies show that IGF-1 can promote the differentiation of adipose-derived stem cells and endothelial cells by the PI3K/AKT signaling pathway⁴². Meanwhile, the IGF-1-induced activation of the AKT signaling pathway also involved the proliferation, migration, osteogenesis, and angiogenesis of hDPSCs²¹. However, the underlying molecular mechanism of proliferation and odontogenic differentiation for hDPSCs needs further research.

In conclusion, we found that exogenous IGF-1 promoted the proliferation of hDPSCs in a concentration-dependent manner in vitro and progressively promoted odontogenic differentiation of hDPSCs in vivo. This suggests that modifying IGF-1 signaling may offer potential strategies for hDPSCs-based tissue engineering to regenerate dental tissues.

Declarations Ethics approval and consent to participate

This study was conducted following the Helsinki Declaration. This study was conducted with approval from the Ethics Committee of the Second Hospital of Hebei Medical University. Written informed consent was obtained from all participants. This study was conducted under the principles of ethical animal research outlined in the Basel Declaration and the ethical guidelines by the International Council for Laboratory Animal Science (ICLAS). This study was conducted following NC3Rs ARRIVE guidelines.

Competing interest

The authors had no separate personal, financial, commercial, or academic conflicts of interest.

Funding

Not applicable.

Authors ORCID'number

- Yan Wang (YW):
0009-0004-4323-0268
- Nan Du (ND):
0009-0009-3942-3872
- Cong-na Liu (CNL):
0009-0002-2022-0329
- Wen-jing Li (WJL):
0009-0007-4067-5845

Author contributions

Conception and design: Wang W; administrative support: YW and ND; provision of study materials or patients: ND and CNL; collection and assembly of data: CNL and WJL; data analysis and interpretation: YW and WJL; manuscript writing: all authors and final approval of manuscript: all authors.

REFERENCES

1. Liu S, Wang S, Dong Y. Evaluation of a bioceramic as a pulp capping agent in vitro and in vivo. *J Endod.* 2015;41(5):652-657. <https://doi.org/10.1016/j.joen.2014.12.009>.
2. Cooper PR, Takahashi Y, Graham LW, Simon S, Imazato S, Smith AJ. Inflammation-regeneration interplay in the dentine-pulp complex. *J Dent* 2010;38(9):687-697. <https://doi.org/10.1016/j.jdent.2010.05.016>.
3. Wang YL, Hu YJ, Zhang FH. Effects of GP-NMB on proliferation and odontoblastic differentiation of human dental pulp cells. *Int J Clin Exp Pathol* 2015;8(6):6498-6504.
4. Miura M, Gronthos S, Zhao M, Lu B, Fisher LW, Robey PG, Shi S. SHED: stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci USA.* 2003;100(10):5807-5812. <https://doi.org/10.1073/pnas.0937635100>.
5. Jo YY, Lee HJ, Kook SY, Choung HW, Park JY, Choung JH, Choung YH, Kim ES, Yang HC, Choung PH. Isolation and

- characterization of postnatal stem cells from human dental tissues. *Tissue Eng.* 2007;13(4):767-773. <https://doi.org/10.1089/ten.2006.0192>.
6. **Paino F, Ricci G, De Rosa A, D'Aquino R, Laino L, Pirozzi G, Tirino V, Papaccio G.** Ecto-mesenchymal stem cells from dental pulp are committed to differentiate into active melanocytes. *Eur Cell Mater.* 2010;20:295-305. <https://doi.org/10.22203/ecm.v020a24>.
 7. **Kim BC, Bae H, Kwon IK, Lee EJ, Park JH, Khademhosseini A, Hwang YS.** Osteoblastic/cementoblastic and neural differentiation of dental stem cells and their applications to tissue engineering and regenerative medicine. *Tissue Eng Part B Rev.* 2012;18(3):235-244. <https://doi.org/10.1089/ten.TEB.2011.0642>.
 8. **Smith JG, Smith AJ, Shelton RM, Cooper PR.** Dental pulp cell behavior in biomimetic environments. *J Dent Res.* 2015;94(11):1552-1559. <https://doi.org/10.1177/0022034515599767>.
 9. **Chen FM, Zhang J, Zhang M, An Y, Chen F, Wu ZF.** A review on endogenous regenerative technology in periodontal regenerative medicine. *Biomaterials.* 2010;31(31):7892-7927. <https://doi.org/10.1016/j.biomaterials.2010.07.019>.
 10. **Liang C, Liao L, Tian W.** Stem cell-based dental pulp regeneration: insights from signaling pathways. *Stem Cell Rev Rep.* 2021;17(4):1251-1263. <https://doi.org/10.1007/s12015-020-10117-3>.
 11. **Suzuki T, Lee CH, Chen M, Zhao W, Fu SY, Qi JJ, Chotkowski G, Eising SB, Wong A, Mao JJ.** Induced migration of dental pulp stem cells for in vivo pulp regeneration. *J Dent Res.* 2011;90(8):1013-1018. <https://doi.org/10.1177/0022034511408426>.
 12. **Huang GT, Gronthos S, Shi S.** Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. *J Dent Res.* 2009;88(9):792-806. <https://doi.org/10.1177/0022034509340867>.
 13. **Pierdomenico L, Bonsi L, Calvitti M, Rondelli D, Arpinati M, Chirumbolo G, Becchetti E, Marchionni C, Alviano F, Fossati V, Staffolani N, Franchina M, Grossi A, Bagnara GP.** Multipotent mesenchymal stem cells with immunosuppressive activity can be easily isolated from dental pulp. *Transplantation.* 2005;80(6):836-842. <https://doi.org/10.1097/01.tp.0000173794.72151.88>.
 14. **Chen FM, Zhang M, Wu ZF.** Toward delivery of multiple growth factors in tissue engineering. *Biomaterials.* 2010;31(24):6279-6308. <https://doi.org/10.1016/j.biomaterials.2010.04.053>.
 15. **Kim SG, Zhou J, Solomon C, Zheng Y, Suzuki T, Chen M, Song S, Jiang N, Cho S, Mao JJ.** Effects of growth factors on dental stem/progenitor cells. *Dent Clin North Am.* 2012;56(3):563-575. <https://doi.org/10.1016/j.cden.2012.05.001>.
 16. **Fang Y, Wang LP, Du FL, Liu WJ, Ren GL.** Effects of insulin-like growth factor I on alveolar bone remodeling in diabetic rats. *J Periodontol Res.* 2013;48(2):144-150. <https://doi.org/10.1111/j.1600-0765.2012.01512.x>.
 17. **Chen FM, Zhao YM, Wu H, Deng ZH, Wang QT, Zhou W, Liu Q, Dong GY, Li K, Wu ZF, Jin Y.** Enhancement of periodontal tissue regeneration by locally controlled delivery of insulin-like growth factor-I from dextran-co-gelatin microspheres. *J Control Release.* 2006;114(2):209-222. <https://doi.org/10.1016/j.jconrel.2006.05.014>.
 18. **Feng X, Huang D, Lu X, Feng G, Xing J, Lu J, Xu K, Xia W, Meng Y, Tao T, Li L, Gu Z.** Insulin-like growth factor 1 can promote proliferation and osteogenic differentiation of human dental pulp stem cells via mTOR pathway. *Dev Growth Differ.* 2014;56(9):615-624. <https://doi.org/10.1111/dgd.12179>.
 19. **Yu Y, Mu J, Fan Z, Lei G, Yan M, Wang S, Tang C, Wang Z, Yu J, Zhang G.** Insulin-like growth factor 1 enhances the proliferation and osteogenic differentiation of human periodontal ligament stem cells via ERK and JNK MAPK pathways. *Histochem Cell Biol.* 2012;137(4):513-525. <https://doi.org/10.1007/s00418-011-0908-x>.
 20. **Lu W, Xu W, Li J, Chen Y, Pan Y, Wu B.** Effects of vascular endothelial growth

- factor and insulin growth factor1 on proliferation, migration, osteogenesis and vascularization of human carious dental pulp stem cells. *Mol Med Rep.* 2019;20(4):3924-3932. <https://doi.org/10.3892/mmr.2019.10606>.
21. Lins-Candeiro CL, Paranhos LR, de Oliveira Neto NF, Ribeiro RAO, de-Souza-Costa CA, Guedes FR, da Silva WHT, Turrioni AP, Santos Filho PCF. Viability and oxidative stress of dental pulp cells after indirect application of chemomechanical agents: An in vitro study. *Int Endod J.* 2023; 57(3):315-327. doi: 10.1111/iej.14013.
 22. Schreier C, Rothmiller S, Scherer MA, Rummel C, Steinritz D, Thiermann H, Schmidt A. Mobilization of human mesenchymal stem cells through different cytokines and growth factors after their immobilization by sulfur mustard. *Toxicol Lett.* 2018;293:105-111. <https://doi.org/10.1016/j.toxlet.2018.02.011>.
 23. Martin-Del-Campo M, Rosales-Ibañez R, Alvarado K, Sampedro JG, Garcia-Sepulveda CA, Deb S, San Román J, Rojo L. Strontium folate loaded biohybrid scaffolds seeded with dental pulp stem cells induce in vivo bone regeneration in critical sized defects. *Biomater Sci.* 2016;4(11):1596-1604. <https://doi.org/10.1039/c6bm00459h>.
 24. Woo SM, Lim HS, Jeong KY, Kim SM, Kim WJ, Jung JY. Vitamin D promotes odontogenic differentiation of human dental pulp cells via ERK activation. *Mol Cells.* 2015;38(7):604-609. <https://doi.org/10.14348/molcells.2015.2318>.
 25. Teng CF, Jeng LB, Shyu WC. Role of insulin-like growth factor 1 receptor signaling in stem cell stemness and therapeutic efficacy. *Cell Transplant.* 2018;27(9):1313-1319. <https://doi.org/10.1177/0963689718779777>.
 26. Werle SB, Lindemann D, Steffens D, Demarco FF, de Araujo FB, Pranke P, Casagrande L. Carious deciduous teeth are a potential source for dental pulp stem cells. *Clin Oral Investig.* 2016;20(1):75-81. <https://doi.org/10.1007/s00784-015-1477-5>.
 27. Gnanasegaran N, Govindasamy V, Abu Kasim NH. Differentiation of stem cells derived from carious teeth into dopaminergic-like cells. *Int Endod J.* 2016;49(10):937-949. <https://doi.org/10.1111/iej.12545>. Epub 2015 Oct 3.
 28. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A.* 2000;97(25):13625-13630. <https://doi.org/10.1073/pnas.240309797>.
 29. Almushayt A, Narayanan K, Zaki AE, George A. Dentin matrix protein 1 induces cytodifferentiation of dental pulp stem cells into odontoblasts. *Gene Ther.* 2006;13(7):611-620. <https://doi.org/10.1038/sj.gt.3302687>.
 30. Zhang W, Walboomers XF, Shi S, Fan M, Jansen JA. Multilineage differentiation potential of stem cells derived from human dental pulp after cryopreservation. *Tissue Eng.* 2006;12(10):2813-2823. <https://doi.org/10.1089/ten.2006.12.2813>.
 31. Yang X, Zhang W, van den Dolder J, Walboomers XF, Bian Z, Fan M, Jansen JA. Multilineage potential of STRO-1+ rat dental pulp cells in vitro. *J Tissue Eng Regen Med.* 2007;1(2):128-135. <https://doi.org/10.1002/term.13>.
 32. Moeller M, Pink C, Endlich N, Endlich K, Grabe HJ, Völzke H, Dörr M, Nauck M, Lerch MM, Köhling R, Holtfreter B, Kocher T, Fuellen G. Mortality is associated with inflammation, anemia, specific diseases and treatments, and molecular markers. *PLoS One.* 2017;12(4):e0175909. <https://doi.org/10.1371/journal.pone.0175909>.
 33. Alarçin E, Lee TY, Karuthedom S, Mohammadi M, Brennan MA, Lee DH, Marrella A, Zhang J, Syla D, Zhang YS, Khademhosseini A, Jang HL. Injectable shear-thinning hydrogels for delivering osteogenic and angiogenic cells and growth factors. *Biomater Sci.* 2018;6(6):1604-1615. <https://doi.org/10.1039/c8bm00293b>.
 34. Shinohara Y, Tsuchiya S, Hatae K, Honda MJ. Effect of vitronectin bound to insulin-like growth factor-I and insulin-like growth factor binding protein-3 on porcine ena-

- mel organ-derived epithelial cells. *Int J Dent*. 2012;2012:386282. <https://doi.org/10.1155/2012/386282>.
35. **Catón J, Bringas P Jr, Zeichner-David M.** Establishment and characterization of an immortal mouse-derived odontoblast-like cell line to evaluate the effect of insulin-like growth factors on odontoblast differentiation. *J Cell Biochem*. 2007;100(2):450-463. <https://doi.org/10.1002/jcb.21053>.
36. **Schminke B, Vom Orde F, Gruber R, Schliephake H, Bürgers R, Miosge N.** The pathology of bone tissue during peri-implantitis. *J Dent Res*. 2015;94(2):354-361. <https://doi.org/10.1177/0022034514559128>.
37. **Onishi T, Kinoshita S, Shintani S, Sobue S, Ooshima T.** Stimulation of proliferation and differentiation of dog dental pulp cells in serum-free culture medium by insulin-like growth factor. *Arch Oral Biol*. 1999;44(4):361-371. [https://doi.org/10.1016/s0003-9969\(99\)00007-2](https://doi.org/10.1016/s0003-9969(99)00007-2).
38. **Wang S, Mu J, Fan Z, Yu Y, Yan M, Lei G, Tang C, Wang Z, Zheng Y, Yu J, Zhang G.** Insulin-like growth factor 1 can promote the osteogenic differentiation and osteogenesis of stem cells from apical papilla. *Stem Cell Res*. 2012;8(3):346-356. <https://doi.org/10.1016/j.scr.2011.12.005>.
39. **Chen C, Xu Y, Song Y.** IGF-1 gene-modified muscle-derived stem cells are resistant to oxidative stress via enhanced activation of IGF-1R/PI3K/AKT signaling and secretion of VEGF. *Mol Cell Biochem*. 2014;386(1-2):167-175. <https://doi.org/10.1007/s111010-013-1855-8>.
40. **Liu C, Zhang Z, Tang H, Jiang Z, You L, Liao Y.** Crosstalk between IGF-1R and other tumor promoting pathways. *Curr Pharm Des*. 2014;20(17):2912-2921. <https://doi.org/10.2174/1381612811319990596>.
41. **Girnitá L, Worrall C, Takahashi S, Se-regard S, Girnitá A.** Something old, something new and something borrowed: emerging paradigm of insulin-like growth factor type 1 receptor (IGF-1R) signaling regulation. *Cell Mol Life Sci*. 2014;71(13):2403-2427. <https://doi.org/10.1007/s00018-013-1514-y>.
42. **Wan Y, Zhuo N, Li Y, Zhao W, Jiang D.** Autophagy promotes osteogenic differentiation of human bone marrow mesenchymal stem cell derived from osteoporotic vertebrae. *Biochem Biophys Res Commun*. 2017;488(1):46-52. <https://doi.org/10.1016/j.bbrc.2017.05.004>.