

Rutin as a potential therapeutic agent for arsenic-induced testicular damage: Antioxidant, anti-inflammatory, and anti-apoptotic effects.

Yunyun Wang^{1,2}, Zengrong Hua^{2,3}, Jin Qing Hui³ and Dandan Qiu⁴

¹Department of Pharmacy, The First Affiliated Hospital of Zhejiang Chinese Medical University, Zhejiang Hospital of Traditional Chinese Medicine, Hangzhou, Zhejiang, China.

²Department of Urology, Huai'an Clinical Medical College of Jiangsu University (Huai'an Hospital of Huai'an City), Huai'an, Jiangsu, China.

³Department of Surgical, Shaanxi Kangfu Hospital, Xian, Shaanxi, China.

⁴Department of Urology, The First Affiliated Hospital of Zhejiang Chinese Medical University, Zhejiang Hospital of Traditional Chinese Medicine, Hangzhou Zhejiang, China.

*Yunyun Wang and Zengrong Hua are co-first authors; they contributed equally to this work.

Keywords: Apoptosis; Arsenic; Inflammation; Rutin, Spermatogenesis; Testicular Toxicity.

Abstract. Arsenic exposure is associated with male reproductive toxicity, including impaired spermatogenesis, reduced sperm quality, and disrupted hormone levels. Rutin, a natural flavonoid, has demonstrated protective effects in various organs owing to its antioxidant, anti-inflammatory, and antiapoptotic properties. This study aimed to assess the efficacy of rutin in ameliorating testicular toxicity caused by sodium arsenite in laboratory rats. Sprague-Dawley rats were randomly assigned to normal, arsenic control, rutin (25, 50, and 100 mg/kg), and Co-enzyme Q10 (10 mg/kg) treatment groups. Testicular damage was induced by oral administration of sodium arsenite (10 mg/kg, for two days). Rutin and Co-enzyme Q10 were administered orally for 15 and seven days, respectively. The serum hormone levels, sperm parameters, testicular mitochondrial enzymes, oxidative stress markers, inflammatory cytokines, apoptotic proteins, and histopathology were assessed. Arsenic exposure significantly decreased ($p < 0.001$) sperm parameters (count, motility, and viability), serum hormone levels (follicle-stimulating hormone, luteinizing hormone, and testosterone), and mitochondrial enzyme activity. Rutin (50 and 100 mg/kg) significantly ($p < 0.01$ and $p < 0.001$) attenuated arsenic-induced alterations in a dose-dependent manner, improving organ weight, sperm parameters, and hormone levels. Rutin also improved mitochondrial complex activity and testicular architecture. In contrast, elevated oxidative stress (reduced glutathione, superoxide dismutase, nitric oxide, and lipid peroxidation), inflammation (tumor necrosis factor-alpha,

interleukin-6, and interleukin-1 β), and apoptosis (caspase-3 and caspase-9 protein expression) were ameliorated by rutin. In conclusion, rutin demonstrated significant protective effects against sodium arsenite-induced testicular toxicity in rats by reducing oxidative stress, inflammation, and apoptosis. These findings suggest that rutin has therapeutic potential in mitigating arsenic-related reproductive toxicity.

Rutina, un agente terapéutico potencial para el daño testicular inducido por arsénico: Efecto antioxidante, antiinflamatorio, y anti-apoptótico.

Invest Clin 2025; 66 (4): 408 – 425

Palabras clave: Apoptosis; Arsénico; Inflamación; Rutina; Espermato-génesis; Toxicidad Testicular.

Resumen. La exposición al arsénico se asocia con toxicidad reproductiva masculina que incluye alteración de la espermato-génesis, reducción de la calidad del esperma y alteración de los niveles hormonales. La rutina, un flavonoide natural, ha demostrado efectos protectores en varios órganos debido a sus propiedades antioxidantes, antiinflamatorias y antiapoptóticas. El propósito del trabajo fue evaluar la eficacia de la rutina para reducir la toxicidad testicular causada por el arsenito de sodio en ratas de laboratorio. Ratas (Sprague-Dawley) fueron asignadas aleatoriamente a grupos de tratamiento normal, control de arsénico, rutina (25, 50 y 100 mg/kg) y coenzima Q10 (10 mg / kg). El daño testicular se indujo mediante la administración oral de arsenito de sodio (10 mg/kg, 2 días). La rutina y la coenzima Q10 se administraron por vía oral durante 15 y 7 días, respectivamente. Se evaluaron los niveles séricos de hormonas, los parámetros de los espermatozoides, las enzimas mitocondriales testiculares, los marcadores de estrés oxidativo, las citocinas inflamatorias, las proteínas apoptóticas y la histopatología. La exposición al arsénico disminuyó significativamente ($p<0,001$) los parámetros espermáticos (recuento, motilidad y viabilidad), y los niveles séricos de hormonas (hormona estimulante del folículo, hormona luteinizante y testosterona) y la actividad de las enzimas mitocondriales. La rutina (50 y 100 mg/kg) atenuó significativamente ($p<0,01$ y $p<0,001$) las alteraciones inducidas por el arsénico de manera dosis-dependiente, mejorando el peso de los órganos, los parámetros espermáticos y los niveles hormonales. La rutina también mejoró la actividad del complejo mitocondrial y la arquitectura testicular, mientras que el estrés oxidativo elevado (glutatión reducido y superóxido dismutasa, óxido nítrico y peroxidación lipídica), la inflamación (factor de necrosis tumoral alfa, interleucina-6 e interleucina-1 β) y la apoptosis (expresión de proteínas caspasa-3 y caspasa-9) mejoraron con la rutina. En conclusión, la rutina demostró efectos protectores significativos contra la toxicidad testicular inducida por el arsenito de sodio en ratas al reducir el estrés oxidativo, la inflamación y la apoptosis. Estos hallazgos sugieren que la rutina tiene potencial terapéutico para mitigar la toxicidad reproductiva asociada al arsénico.

Received: 12-08-2025 Accepted: 22-09-2025

INTRODUCTION

Arsenic is a metalloid commonly found in the environment, known for its toxicity and important public health effects ^{1,2}. It mainly impacts populations through contaminated drinking water, often from natural sources and human activities such as industrial discharge and the use of arsenic-based pesticides ^{3,4}. According to the World Health Organization (WHO), more than 140 million people worldwide are exposed to arsenic-contaminated water exceeding the safe limit of 10 µg/L ⁵. Furthermore, long-term exposure to arsenic has been associated with numerous health problems, including cancer and various non-cancerous conditions like cardiovascular diseases, diabetes, and reproductive issues ^{6,7}. Sodium arsenite is recognized for causing testicular toxicity through mechanisms such as oxidative stress, apoptosis, and inflammation. Researchers believe that inflammation, oxidative/nitrosative stress, and apoptosis are key factors in arsenic-induced testicular damage ^{6,8}.

The impact of arsenic toxicity on reproductive health has become an increasing concern, particularly regarding the male reproductive system ⁹. Arsenic exposure is linked to various reproductive problems, including male infertility, decreased sperm quality, and disruptions in testosterone production. This results in reduced weights of reproductive organs, more sperm abnormalities, and apoptosis in testicular cells ¹⁰. The harmful effects of arsenic on male fertility involve several mechanisms, such as interference with steroidogenesis and mitochondrial dysfunction in reproductive tissues ⁹. These disruptions lead to lower testosterone levels, decreased sperm count, and abnormal sperm morphology ¹¹. Additionally, changes in spermatogenesis and reduced gonadotropin levels have been observed, further emphasizing the reproductive risks of arsenic ³. Moreover, arsenic exposure causes increased production of reactive oxygen species (ROS), indicating the role of oxidative

stress in its genotoxic effects ¹². Researchers have documented that testicular toxicity caused by sodium arsenite is partly due to its ability to induce apoptosis through the p53 pathway, as shown in zebrafish studies where sodium arsenite exposure elevated apoptosis markers and decreased global DNA methylation ¹³. Furthermore, experiments with *Cae-norhabditis elegans* demonstrated that sodium arsenite can cause cell cycle arrest and germline apoptosis, both of which depend on concentration and exposure time ¹².

Several protective agents have been studied to reduce testicular toxicity caused by sodium arsenite. Melatonin has been shown to decrease arsenic-induced testicular cell death, oxidative stress, and tissue damage by boosting antioxidant enzyme activity and lowering lipid peroxidation ¹⁴. Similarly, *Salvia hispanica* (chia seeds) proved effective in decreasing testicular toxicity by improving sperm quality, serum sex hormone levels, and antioxidant enzyme activity, thanks to its flavonoid content and antioxidant properties ¹⁵. Hesperidin and lipoic acid were also tested for their protective effects against liver, kidney, and testicular toxicity when administered with sodium arsenite ¹⁶. Rutin is a natural flavonoid with a wide range of pharmacological benefits, including antioxidant, antimicrobial, antifungal, anti-allergic, anti-cancer, anti-inflammatory, and antiapoptotic effects ^{4,17-22}. Additionally, rutin has been shown to help manage neurodegenerative diseases and metabolic conditions like diabetes due to its cell-protective effects ¹⁹. Its protective properties have been documented in the liver, kidneys, and heart due to its anti-inflammatory, antioxidant, and antiapoptotic actions, making it a candidate for protecting these organs from toxic agents ⁴. However, the impact of rutin on testicular toxicity caused by sodium arsenite has not yet been studied. Given its protective effects on other organs, rutin might provide similar benefits in reducing testicular damage from sodium arsenite. This study examined the effectiveness of rutin in alleviating testicular toxicity caused by sodium arsenite in male rats.

MATERIALS AND METHODS

Experimental design

Rats (male, Sprague-Dawley, weighing 200-220 g, sourced from the animal house of Shaanxi Kangfu Hospital, China) were randomly divided into the following groups (n=6 per group): normal, arsenic control (As), rutin (Sigma Chemical Co., St Louis, Missouri, United States; 25, 50, and 100 mg/kg), and Co-enzyme Q₁₀ (Medicines Pvt. Ltd. Mumbai, India; 10 mg/kg). Testicular damage was induced in rats (except the normal group) by oral administration of sodium arsenite (Otto Chemicals, India; 10 mg/kg, for two days)^{23,24}. Rats in the arsenic control and normal groups were treated with double-distilled water (10 mL/kg). Three different dosages of rutin (25, 50, and 100 mg/kg) were selected based on previous studies^{21,22} and were administered orally for 15 days. Co-enzyme Q₁₀ was administered for seven consecutive days before arsenite administration and continued for up to 15 days. On the 16th day, the rats were put under anesthesia using ether, and blood was drawn through retro-orbital puncture. Each blood sample was placed in a separate vial for analysis of serum parameters. Serum follicle-stimulating hormone, luteinizing hormone, and testosterone levels were quantified according to the manufacturer's instructions for the rat-specific ELISA kit (Thermo Scientific, Rockford, IL, USA). The rats were euthanized by carbon dioxide asphyxiation, and the testes were rapidly removed and stored at -80°C for biochemical parameters. Other organs, such as the epididymis and prostate, were isolated and weighed. The epididymal sperm count and motility were determined according to a previously reported method²⁵.

Biochemical estimation of testis homogenate

Supernatant of the tissue homogenate was employed to estimate lipid peroxidation [malondialdehyde (MDA) content], nitric oxide (NO content), reduced glutathione

(GSH), and superoxide dismutase (SOD)] as described previously²⁶⁻²⁹. Testicular mitochondrial enzyme activities, including Complex I (nicotinamide-adenine dinucleotide (NADH) dehydrogenase activity), Complex II (succinate dehydrogenase (SDH) activity), Complex III (mitochondrial redox activity), and Complex IV (cytochrome oxidase assay) were estimated according to previously reported methods^{30,31}.

Testicular interleukins (interleukins (ILs); IL-6 (ERA31RB) and IL-1 β (BMS630)) and tumor necrosis factor-alpha (TNF- α ; ERA56RB) were quantified in the testis homogenate using Enzyme-linked immunosorbent assay (ELISA) kits (Thermo Scientific, Rockford, IL, USA). Briefly, 500 mg of testis tissue samples were homogenized with a mechanical homogenizer in 5 ml of phosphate-buffered saline at 3000 rpm. The homogenate was centrifuged for 30 min at 20,000 rpm (4°C) in a cryo centrifuge (Eppendorf), and the supernatant was used to determine ILs and TNF- α . Briefly, the quantification of ILs and TNF- α was performed using the Thermo Scientific Rat ILs and TNF- α immunoassay kit, following the instructions provided. The Rat ILs and TNF- α immunoassay was a 4.5 h solid phase designed to measure rat ILs and TNF- α levels. The assay employed a sandwich enzyme immunoassay principle. A monoclonal antibody specific for rat ILs and TNF- α was pre-coated on the microplates. Standards, control, and samples were pipetted into the wells, and the immobilized antibody thus bound any rat ILs or TNF- α present in the sample. After washing away the unbound substance, an enzyme-linked polyclonal antibody specific for rat ILs or TNF- α was pipetted into the microtitre wells. Any unbound antibody was washed off, and then a substrate solution was added to the wells. The enzymatic reaction produced a blue product that turned yellow upon addition of the stop solution. The intensity of the generated color was measured and was proportional to the amount of rat ILs or TNF- α bound in the initial steps. A standard curve was run

on each assay plate using recombinant ILs or TNF- α in serial dilutions. The sample values were then read and calculations made according to the standard curve. Values were expressed as means \pm S.E.M. The levels of ILs or TNF- α were expressed as units per mg of gastric tissue.

The protein expression of caspase-3 and caspase-9 was assessed by western blot in the testis³²⁻³⁴. Briefly, the testis was sonicated in Tissue Protein Extraction Reagent (Thermo Fisher Scientific, Inc., Mumbai, Maharashtra, India). The lysates were centrifuged at 10,000 \times g for 10 min at 4°C. Protein concentration was determined using a Bicinchoninic Acid (BCA) assay kit (Beyotime Shanghai, China) on ice for 30 min. Equal amounts of extracted protein samples (50 μ g) were separated by 10% SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis) and transferred onto polyvinylidene difluoride membranes. The membranes were blocked with 5% non-fat dry milk at 37°C for 1 hr and incubated overnight at 4°C with the primary antibodies recognizing caspase-3 (ab4051; 1/200 dilution; 31 kDa) and caspase-9 (ab202068; 1/2000 dilution; 46 kDa; Abcam, Cambridge, MA, USA). In addition, an anti-rabbit horseradish-linked secondary antibody (goat anti-rabbit IgG H&L; ab97051) was used, which was incubated at 37°C for 2 hr. Protein bands were visualized using the Chemiluminescent kit (Bio-Rad Laboratories, Inc., Mumbai, Maharashtra, India), and glyceraldehyde 3-phosphate dehydrogenase (GAPDH; EPR6256, ab128915; 1/10000 dilution; 36 kDa) served as the loading control.

Histopathological analysis of testis

Other testis samples were preserved in 10% formalin for 24 hours. These samples underwent dehydration and were immersed in xylene for one hour, repeated three times, followed by treatment with ethyl alcohol at concentrations of 70%, 90%, and 100% for two hours. The infiltration and impregnation process involved treating the samples with

paraffin wax twice, with each session lasting 1 hour. The tissue samples were then sliced into sections with a thickness of 3-5 μ m and stained using hematoxylin and eosin (H&E). The sections were mounted on slides with Distrene Phthalate Xylene (DPX) serving as the mounting medium. A light microscope was used to examine the sections for histopathological characteristics and cell infiltration. The observed changes in histological features were categorized into grades ranging from 0 to 4 according to a previously reported method³⁵.

Statistical analysis

The data are presented as mean \pm standard error of the mean (SEM). GraphPad Prism 5.0 software (GraphPad, San Diego, CA, USA) was utilized for data analysis. Biochemical parameter data were examined using one-way analysis of variance, followed by Tukey's multiple range test for parametric results, while the Kruskal-Wallis test was used for non-parametric outcomes. A p of less than 0.05 was deemed statistically significant. Correlation coefficients were calculated using a two-sided Fisher's test.

RESULTS

Attenuation of arsenic-induced alteration in body weight and organ weights by rutin

Table 1 presents the descriptive results for various body and organ weight parameters across the different treatment groups. The arsenic (As) control group had a significant reduction ($p<0.001$) in organ weights (testes, epididymis, and prostate) and body weight compared with the normal group. CoQ treatment resulted in a significant improvement ($p<0.001$) in body weight and organ weights compared to the arsenic control group. Rutin (50 and 100 mg/kg) administration also effectively ($p<0.01$ and $p<0.001$) and dose-dependently increased the body, testes, epididymis, and prostate weights relative to the arsenic-exposed group.

Table 1. Attenuation of arsenic-induced alteration in body weight, organ weights (testes, epididymis, and prostate), and sperm parameters (count, motility, and viability) by rutin.

| Parameters | Normal | As Control | CoQ (10) | R (25) | R (50) | R (100) |
|--|----------------|------------------|------------------|----------------|------------------|-------------------|
| Body weight (gm) | 281.8 ± 2.86 | 217.50 ± 3.73### | 275.70 ± 2.69*** | 214.70 ± 2.80 | 247.00 ± 4.02** | 262.20 ± 2.77*** |
| Testes weight (gm) | 1.79 ± 0.04 | 1.03 ± 0.07### | 1.59 ± 0.04*** | 1.07 ± 0.05 | 1.51 ± 0.06*** | 1.68 ± 0.02*** |
| Testes weight/Body weight ($\times 10^{-3}$) | 6.34 ± 0.15 | 4.75 ± 0.39### | 5.76 ± 0.18*** | 4.99 ± 0.21 | 6.13 ± 0.33** | 6.41 ± 0.12*** |
| Epididymis weight (gm) | 0.51 ± 0.01 | 0.19 ± 0.02### | 0.45 ± 0.02*** | 0.23 ± 0.01 | 0.33 ± 0.01** | 0.45 ± 0.01 *** |
| Epididymis weight/Body weight ($\times 10^{-3}$) | 1.80 ± 0.03 | 0.87 ± 0.07### | 1.62 ± 0.07*** | 1.09 ± 0.07 | 1.32 ± 0.04** | 1.73 ± 0.04*** |
| Prostate weight (mg) | 612.30 ± 10.94 | 308.30 ± 9.78### | 560.70 ± 9.02*** | 298.70 ± 11.44 | 478.30 ± 11.78** | 541.70 ± 11.45*** |
| Prostate weight/Body weight ($\times 10^{-3}$) | 2.18 ± 0.04 | 1.42 ± 0.05### | 2.03 ± 0.04*** | 1.39 ± 0.04 | 1.94 ± 0.07** | 2.07 ± 0.05*** |
| Sperm count (millions/mL) | 60.17 ± 1.25 | 34.00 ± 0.97### | 56.83 ± 1.30*** | 33.83 ± 0.75 | 45.00 ± 1.21** | 51.83 ± 1.25*** |
| Sperm motility (%) | 71.33 ± 1.05 | 40.33 ± 1.02### | 62.00 ± 1.00*** | 44.33 ± 0.95 | 49.17 ± 1.49** | 59.50 ± 0.67*** |
| Dead sperm (%) | 28.67 ± 1.05 | 59.67 ± 1.02### | 38.00 ± 1.00*** | 55.67 ± 0.95 | 50.83 ± 1.49** | 40.50 ± 0.67*** |
| Abnormal sperm (%) | 12.17 ± 0.60 | 36.50 ± 0.43### | 17.17 ± 0.54*** | 34.17 ± 0.48 | 24.67 ± 0.42** | 19.33 ± 0.76*** |

Data analysis: One-way ANOVA (post-hoc test: Tukey's multiple range test). Data are reported as mean ± SEM (n=6). Statistically significant compared with $###$ normal rats, ** and *** as control rats. $###$ $p < 0.001$, ** $p < 0.01$ and *** $p < 0.001$. Arsenic (As), Co-enzyme Q₁₀ (CoQ (10)) and Rutin (R).

Table 2. Attenuation of arsenic-induced alteration in serum luteinizing hormone, follicle stimulating hormone, testosterone levels and testicular mitochondrial complex activities by rutin.

| Parameters | Normal | As Control | CoQ (10) | R (25) | R (50) | R (100) |
|---|------------------|---------------------|---------------------|------------------|--------------------|---------------------|
| Luteinizing hormone (ng/mL) | 1.88 ± 0.04 | 0.26 ± 0.05### | 0.33 ± 0.03 | 0.36 ± 0.04 | 0.71 ± 0.05** | 1.21 ± 0.04*** |
| Follicle stimulating hormone (ng/mL) | 1.37 ± 0.02 | 0.16 ± 0.02### | 0.15 ± 0.02 | 0.26 ± 0.02 | 0.64 ± 0.02** | 0.92 ± 0.02*** |
| Testosterone (ng/mL) | 3.69 ± 0.04 | 1.41 ± 0.05### | 1.32 ± 0.06 | 1.5 ± 0.06 | 2.39 ± 0.04** | 2.86 ± 0.05*** |
| Testicular parameters | | | | | | |
| Complex I (nmole of NADH oxidized/min/mg protein) | 33.36 ± 1.66 | 7.41 ± 1.80### | 30.21 ± 1.79*** | 10.07 ± 1.35 | 16.91 ± 1.25** | 28.31 ± 2.10*** |
| Complex II (nmole/mg protein) | 14.35 ± 0.82 | 4.38 ± 0.78### | 11.86 ± 0.84*** | 4.74 ± 0.62 | 9.42 ± 0.67** | 11.24 ± 0.58*** |
| MTT assay (OD at 540 nm) | 0.45 ± 0.03 | 0.17 ± 0.03### | 0.45 ± 0.03*** | 0.21 ± 0.02 | 0.32 ± 0.01** | 0.39 ± 0.02*** |
| Complex-IV (nmol cyto-C oxidized/min/mg protein) | 6319.00 ± 280.30 | 1008.00 ± 212.70### | 5293.00 ± 254.70*** | 1586.00 ± 191.90 | 3224.00 ± 201.30** | 4515.00 ± 342.00*** |

Data analysis: One-way ANOVA (post-hoc test: Tukey's multiple range test). Data are reported as mean ± SEM (n=6). Statistically significant compared with $###$ normal rats, ** and *** as control rats. $###$ $p < 0.001$, ** $p < 0.01$ and *** $p < 0.001$. Arsenic (As), Co-enzyme Q₁₀ (CoQ (10)), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and Rutin (R).

Attenuation of arsenic-induced alteration in sperm parameters by rutin

The arsenic control group showed a significant reduction ($p<0.001$) in cell count and motility compared to the control group. However, the percentage of dead and abnormal sperm was markedly higher ($p<0.001$) in the arsenic control group than that in the normal group. Treatment with CoQ resulted in significant improvements ($p<0.001$) in sperm count and motility compared with the arsenic control group. Furthermore, CoQ administration resulted in a more effective ($p<0.001$) recovery of the percentage of dead and abnormal sperm than arsenic control. Rutin (50 and 100 mg/kg) treatment demonstrated a marked ($p<0.01$ and $p<0.001$) and dose-dependent improvement in sperm parameters relative to the arsenic-exposed group (Table 1).

Attenuation of arsenic-induced alteration in serum follicle-stimulating hormone, luteinizing hormone, and testosterone levels by rutin

Exposure to sodium arsenite significantly decreased ($p<0.001$) the levels of serum follicle-stimulating hormone, luteinizing hormone, and testosterone in the arsenic control group compared with the normal group. Treatment with CoQ and rutin (25 mg/kg) had a minimal impact on serum luteinizing hormone, follicle-stimulating hormone, and testosterone hormone levels. However, rutin (50 and 100 mg/kg) treatments demonstrated effective, dose-dependent increases in all three hormones ($p<0.01$ and $p<0.001$, respectively) compared with the arsenic control group (Table 2).

Attenuation of arsenic-induced alteration in testicular mitochondrial complex activities by rutin

Arsenic exposure led to a significant reduction ($p<0.001$) in mitochondrial complex activity compared to the normal group.

The results presented in Table 2 demonstrate that both CoQ and rutin (50 and 100 mg/kg) significantly improved mitochondrial function in arsenic-exposed rats. Specifically, rutin (50 and 100 mg/kg) administration led to notable ($p<0.01$ and $p<0.001$) and dose-dependent increases in mitochondrial complex activities compared with the arsenic control group. However, rutin (25 mg/kg) did not show any significant potential to improve mitochondrial function or mitigate arsenic-induced damage.

Attenuation of arsenic-induced alteration in testicular oxido-nitrosative stress by rutin

The results presented in Table 3 show the effects of rutin treatment on testicular oxido-nitrosative stress. Arsenic exposure caused a substantial ($p<0.001$) decrease in GSH and SOD levels and a marked ($p<0.001$) elevation in nitric oxide and MDA levels in the arsenic control group compared to the normal group. CoQ treatment significantly ($p<0.001$) improved GSH and SOD levels and effectively reduced ($p<0.001$) nitric oxide and MDA levels compared with the arsenic control group. Rutin (25 mg/kg) treatment showed minimal effects in attenuating arsenic-induced elevated testicular oxido-nitrosative stress compared with the arsenic control group. However, rutin (50 and 100 mg/kg) demonstrated marked ($p<0.01$ and $p<0.001$) and dose-dependent protective effects, reflected by restored antioxidant status (GSH and SOD levels) and reduced oxidative stress markers (nitric oxide and MDA) to normal levels (Table 3).

Attenuation of arsenic-induced alteration in testicular inflammatory markers activities by rutin

The results in Table 3 show significant differences in inflammatory cytokine levels between the normal and arsenic control groups. The arsenic control group exhibited markedly elevated ($p<0.001$) levels of ILs (IL-6 and IL-1 β) and TNF- α compared with

Table 3. Attenuation of arsenic-induced alteration in testicular oxido-nitrosative stress and inflammatory markers by rutin.

| Parameters | Normal | As Control | CoQ (10) | R (25) | R (50) | R (100) |
|------------------------|---------------|-------------------|-------------------|----------------|------------------|-------------------|
| SOD (U/mg of protein) | 13.20 ± 1.10 | 5.19 ± 0.93### | 10.66 ± 0.87*** | 5.45 ± 0.61 | 8.91 ± 0.93** | 10.66 ± 0.96*** |
| GSH (μg/mg of protein) | 13.76 ± 0.64 | 3.90 ± 0.62### | 11.04 ± 0.59*** | 5.36 ± 0.50 | 8.31 ± 0.76** | 10.43 ± 0.56*** |
| MDA (nM/mg of protein) | 0.46 ± 0.10 | 3.14 ± 0.18### | 1.97 ± 0.18*** | 3.00 ± 0.27 | 2.11 ± 0.04** | 1.59 ± 0.19*** |
| NO (μg/mg of protein) | 119.00 ± 9.08 | 284.00 ± 13.96### | 143.90 ± 9.97*** | 261.40 ± 13.64 | 222.10 ± 12.14** | 167.90 ± 9.82*** |
| TNF-α (pg/mL) | 157.00 ± 8.15 | 409.70 ± 12.69### | 222.80 ± 14.07*** | 377.20 ± 14.06 | 330.30 ± 8.02** | 253.70 ± 12.06*** |
| IL-1β (pg/mL) | 17.28 ± 3.86 | 72.12 ± 1.85### | 30.57 ± 2.76*** | 68.46 ± 1.62 | 48.95 ± 2.70** | 42.78 ± 2.41*** |
| IL-6 (pg/mL) | 40.50 ± 5.53 | 152.60 ± 5.18### | 59.41 ± 9.32*** | 155.50 ± 6.21 | 130.40 ± 7.87** | 71.78 ± 7.13*** |

Data analysis: One-way ANOVA (post-hoc test: Tukey's multiple range test). Data are reported as mean ± SEM (n=6). Statistically significant compared with ###normal rats, ** and ***As control rats. ###p < 0.001, **p < 0.01 and ***p < 0.001. Arsenic (As), Co-enzyme Q₁₀ (CoQ (10)), Glutathione (GSH), Interleukin-1 beta (IL-1β), Interleukin-6 (IL-6), Malondialdehyde (MDA), Nitric Oxide (NO), Rutin (R), Superoxide Dismutase (SOD), and Tumor Necrosis Factor-alpha (TNF-α).

the normal group. CoQ treatment substantially (p<0.001) lowered ILs (IL-6 and IL-1β) and TNF-α levels compared to the arsenic control group. Rutin (50 and 100 mg/kg) administration resulted in effective (p<0.01 and p<0.001) and dose-dependent reductions in cytokine levels compared with the arsenic control group. However, a lower dose of rutin (25 mg/kg) did not reduce cytokine levels to levels comparable to those of the arsenic control group, indicating failure to ameliorate arsenic-induced inflammation (Table 3).

Attenuation of arsenic-induced alteration in testicular caspase-3 and caspase-9 protein expressions by rutin

Fig. 1 illustrates the impact of arsenic exposure on markers of apoptosis in the testes, along with their correlation with sperm count. The caspase-3 and caspase-9 protein expression was significantly increased (p<0.001) in the arsenic control group compared with that in the normal group. The substantial elevation in caspase-3 and caspase-9 protein expression was markedly downregulated (p<0.001) by CoQ treat-

ment compared to the arsenic control group. Treatment with rutin (50 and 100 mg/kg) significantly (p<0.01 and p<0.001) and dose-dependently reduced both caspase-3 and caspase-9 relative densities compared to the arsenic control (Figs. 1A and 1B). Furthermore, Figs. 1C and 1D reveal a strong inverse correlation between sperm count and the relative densities of caspase-3 ($R^2 = 0.8168$, p<0.001) and Caspase-9 ($R^2 = 0.7075$, p<0.01), respectively, indicating that lower sperm counts are associated with increased apoptotic activity.

Attenuation of arsenic-induced alteration in testicular histopathology by rutin

Fig. 2 illustrates the histopathological changes in rat testicular tissue after exposure to arsenic and their amelioration across different treatment groups. In the normal group (Fig. 2A), testicular architecture appeared normal, with intact seminiferous tubules containing organized spermatogenic cells and healthy Leydig cells, and no evidence of necrosis. However, it showed mild inflammation (indicated by black arrows). In contrast, the arsenic con-




Fig. 1. Attenuation of arsenic-induced alterations in testicular caspase-3 (A) and caspase-9 (B) protein expression by rutin. Correlation of sperm count with caspase-3 (C) and caspase-9 (D) protein expression. Data analysis: One-way ANOVA (post-hoc test: Tukey's multiple range test). Data are reported as mean \pm SEM ($n=6$). Statistically significant compared with $^{###}$ normal rats, ** and *** As control rats. $^{###}p < 0.001$, $^{**}p < 0.01$ and $^{***}p < 0.001$. Correlation coefficients were determined using a two-sided Fisher's test. Arsenic (As), Co-enzyme Q₁₀ (CoQ (10)), Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH), and Rutin (R). Inflammatory infiltration (black arrow), necrosis (red arrow), and Leydig cells damage (blue arrow).

tral group (Fig. 2B) demonstrated histological aberrations characterized by severe inflammatory infiltration (black arrows), necrosis (red arrows), vacuolated spermatogenic cells (blue arrows), and Leydig cell damage, indicating arsenic-induced toxicity. The quantitative histopathological score in Fig. 2G supports these visual observations, as the arsenic control group exhibited significantly ($p < 0.001$) elevated scores for

inflammatory infiltration, necrosis, Leydig cell damage, and vacuolated spermatogenic cells than the normal group. Treatment with CoQ significantly reduced ($p < 0.001$) these histopathological aberrations compared to the arsenic control group (Fig. 2C), suggesting a protective effect against arsenic-induced testicular injury. The rutin (25 mg/kg) group showed a modest reduction in these scores, but this was not significantly




Fig. 2. Attenuation of arsenic-induced alterations in testicular histopathology by rutin treatment. Representative images of testes from each group (A-F). Quantitative analysis of arsenic-induced alterations in testicular histopathology and its attenuation by rutin (G). Data analysis: One-way ANOVA (post-hoc test: Kruskal-Wallis test). Data are reported as mean \pm SEM (n=6). Statistically significant compared with ***normal rats, ** and ***As control rats. ***p < 0.001, **p < 0.01 and ***p < 0.001. Arsenic (As), Co-enzyme Q₁₀ (CoQ (10)), Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH), and Rutin (R).

different from the higher rutin or CoQ doses (Fig. 2D). Treatment with rutin (50 and 100 mg/kg) showed a significant ($p<0.01$ and $p<0.001$) reduction in arsenic-induced histological aberrations in the testes, reflected in a decrease in inflammatory infiltration, necrosis, Leydig cell damage, and vacuolated spermatogenic cells compared to the arsenic control group (Figs 2E, 2F, and 2G).

DISCUSSION

Arsenic exposure has extensive human health effects, with systemic toxicity affecting multiple organs. It is particularly notorious for its carcinogenic properties, causing skin, lung, and bladder cancers, among others⁶. However, it is crucial to evaluate its adverse effects on male reproductive health as it directly affects male fertility. Arsenic exposure impairs spermatogenesis and reduces sperm quality through oxidative stress and interference with crucial hormonal signaling pathways¹¹. Researchers have explored the protective efficacy of phytonutrients, such as lutein and α -lipoic acid, against arsenic-induced oxidative damage in testicular tissues³⁶. In the present study, rutin, an herbal intervention, was evaluated for its effects on arsenite-induced testicular damage, and the findings suggested that rutin ameliorated testicular toxicity through its anti-apoptotic, anti-inflammatory, and mitochondrial-protective effects in experimental rats.

Integrative proteomic and metabolomic analyses have shown that arsenic exposure significantly alters the proteome and metabolome in rat testes, affecting spermatogenesis and fertilization through disrupted signaling pathways³⁷. Sodium arsenite causes significant testicular damage primarily through oxidative stress, apoptosis, and inflammatory pathways. Studies have demonstrated that chronic exposure to sodium arsenite can lead to a significant decrease in both absolute and relative testicular weight³⁸. This reduction in testicular weight is attributed to the toxic effects

of arsenic on testicular tissue and its interference with normal spermatogenesis and enzyme activities, including decreased acid phosphatase, sorbitol dehydrogenase, and 17 β -hydroxy-steroid dehydrogenase activities, while increasing lactate dehydrogenase and gamma-glutamyl transpeptidase activities. Additionally, arsenite exposure leads to a notable increase in abnormal sperm forms, along with a decrease in sperm count and motility³⁸. Furthermore, arsenic is known to accumulate significantly in reproductive tissues, underscoring its potential to cause prolonged toxic effects³⁸. This accumulation in the testes, epididymis, and prostate is linked to oxidative stress and histopathological alterations, indicating that arsenic's effects are more profound at the cellular level^{8,14}. In the present study, administration of rutin significantly attenuated arsenite-induced decreases in testes, epididymis, and prostate weights, suggesting its protective efficacy in reproductive organs.

Sodium arsenite led to decreased levels of serum testosterone, luteinizing hormone, and follicle-stimulating hormone, and to significant changes in sperm parameters, including reduced sperm count and motility, and an increase in abnormal sperm¹⁵. Serum hormone levels, such as testosterone, FSH, and LH, play crucial roles in regulating male reproductive function³⁹. Testosterone, which is primarily produced in the testes, is vital for normal male reproductive functions and secondary sexual characteristics. FSH and LH, secreted by the pituitary gland, regulate testicular function, including steroidogenesis and spermatogenesis⁴⁰. Chronic exposure to arsenite is often associated with reduced testosterone levels owing to impaired testicular function. This can be attributed to the direct cytotoxic effects of arsenite on Leydig cells, which are responsible for testosterone production. Additionally, alterations in hormone levels can disrupt the hypothalamic-pituitary-gonadal axis, leading to compensatory changes in LH and FSH levels. In this study, arsenite-induced testicular

damage was reflected in changes in serum testosterone, FSH, and LH levels. However, administration of rutin restored the diminished levels of these hormones in the serum.

Oxidative stress plays a crucial role in arsenic-induced toxicity, impacting various biological systems and leading to significant health issues. Arsenic, a toxic metalloid, can cross cellular barriers and accumulate in tissues, contributing to oxidative stress by generating ROS^{41,42}. ROS generation is central to the pathogenesis of arsenic toxicity, as it leads to mitochondrial dysfunction, a critical factor in neurotoxicity, and other health issues^{43,44}. Arsenic-induced oxidative stress leads to mitochondrial damage by disrupting the normal balance between antioxidants and pro-oxidants within cells. This disruption impairs mitochondrial membrane potential, promotes cytochrome c release, and triggers apoptosis through caspase activation⁴². On a systemic level, oxidative stress also causes chromosomal instability and DNA damage, leading to further complications, such as carcinogenesis and neurological disorders⁴⁴⁻⁴⁷. Moreover, arsenic exposure has been shown to deplete antioxidant capacities and elevate oxidative damage biomarkers, such as malondialdehyde (MDA) and nitric oxide, owing to the heightened oxidative environment^{48,49}. These changes underline the oxidative stress mechanism as a cornerstone of arsenic toxicity, which affects various systems, including the reproductive system⁵⁰. Therapeutic strategies to combat arsenic-induced oxidative stress focus on enhancing mitochondrial function and increasing antioxidant defense. Acetyl-L-carnitine (ALC), for instance, has been shown to counteract arsenic-induced oxidative stress by improving antioxidant mechanisms and mitochondrial function, thus offering a potential therapeutic pathway⁴². Other antioxidants, such as curcumin and apigenin, also exhibit protective effects by reducing ROS and supporting cellular antioxidant systems, highlighting their possible roles in mitigating arsenic toxicity^{51,52}. In the present investigation, arsenic-induced toxic-

ity, associated with elevated oxidative stress, contributed to various cellular and systemic damage, as reflected by diminished testicular mitochondrial enzyme activity. However, rutin treatment effectively increased testicular glutathione and superoxide levels and reduced malondialdehyde and nitric oxide levels, suggesting its protective effects against arsenic-induced mitochondrial dysfunction, potentially by mitigating oxidative stress and enhancing mitochondrial energy production.

Inflammation is a significant contributor to arsenic-induced testicular toxicity. Pro-inflammatory cytokines, such as TNF- α and interleukins, mediate this process. TNF- α induces apoptosis and promotes inflammatory responses⁵³. Inflammation induced by TNF- α can activate various signaling pathways, culminating in oxidative stress and tissue damage⁵⁴⁻⁵⁶. It has been observed that arsenic exposure results in the upregulation of inflammatory mediators, such as TNF- α and interleukins, contributing to testicular damage⁵⁷. Furthermore, Caspases 3 and 9 play crucial roles in the apoptotic pathways underlying arsenic-induced testicular toxicity. Caspase-9 is part of the intrinsic or mitochondrial apoptosis pathway, which is often activated by cellular stressors, including toxins such as arsenic. It is responsible for activating downstream effector caspases, including caspase-3^{58,59}. Caspase-3 then executes apoptosis by cleaving cellular substrates, leading to systematic breakdown of cellular components and cell death⁶⁰⁻⁶². In arsenic-induced testicular toxicity, activation of these caspases is indicative of enhanced apoptotic activity, contributing to the observed tissue damage and dysfunction⁵⁷. In the current study, arsenic-induced elevated inflammation and apoptosis, mediated by cytokines such as TNF- α and enzymes such as caspases 3 and 9, highlighted the development of arsenic-induced testicular toxicity. However, rutin treatment mitigated the toxic effects induced by chronic exposure to arsenic via

the amelioration of these pathways, suggesting its therapeutic potential against arsenic-induced testicular toxicity.

This study has some limitations. First, the study focused on acute arsenic exposure (two consecutive days), which may not reflect the effects of chronic, long-term exposure often observed in human populations. Second, the study only used sodium arsenite, whereas environmental arsenic exposure often involves multiple arsenic species. Third, this study did not address the bioavailability and metabolism of rutin in rats, which could affect its efficacy in humans. Lastly, the study focused on rutin alone and did not explore its potential synergistic effects with other protective agents.

As a conclusion, rutin demonstrated significant protective effects against sodium arsenite-induced testicular toxicity in rats. Rutin ameliorated arsenic-induced damage by reducing oxidative stress, inflammation, and apoptosis in testicular tissue, while improving sperm parameters and hormone levels. These findings suggest that rutin may have therapeutic potential in mitigating arsenic-related reproductive toxicity, although further research is needed to elucidate its mechanisms of action and clinical applicability fully.

Acknowledgement

Medical writing support for the development of this manuscript, under the direction of the authors, was provided by Yonnova Scientific Consultancy Pvt. Ltd. in accordance with the Good Publication Practice guidelines.

Funding

This study was funded by the Zhejiang Province Medical and Health Science and Technology Plan Project (2021RC096) and the Zhejiang Traditional Chinese Medicine Science and Technology Plan Project (2023ZL370).

Conflicts of interest

The authors declare that they have no conflict of interest regarding this study.

Availability of data

The datasets generated and/or analyzed during the current study are not publicly available due to confidentiality agreements; supporting data can only be made available to bona fide researcher's subject to a non-disclosure agreement.

Ethical approval

All experiments were performed between 09:00 and 17:00. The experimental protocol was approved by the Institutional Animal Ethics Committee of the Shaanxi Kangfu Hospital (approval no. sxkf2025). All procedures involving animals were conducted in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals.

ORCID number of authors

- Yunyun Wang (YW): 0009-0007-3462-5323
- Zengrong Hua (ZH): 0009-0008-8248-8148
- Jin Qing Hui (JQH): 0009-0008-5583-9213
- Dandan Qiu (DQ): 0009-0009-1376-7853

Author contributions

Each author has made significant contributions to the development of this manuscript. YW: conceived and designed the evaluation and drafted the manuscript; ZH and DQ: performed data acquisition; YW: participated in developing the assessment, performed parts of the statistical analysis, and helped to draft the manuscript; JQH: revised the manuscript and performed the statistical analysis. All the authors have read and approved the final version of the manuscript.

REFERENCES

1. Adil M, Kandhare AD, Ghosh P, Bodhankar SL. Sodium arsenite-induced myocardial bruise in rats: Ameliorative effect of naringin via TGF- β /Smad and Nrf/HO pathways. *Chem Biol Interact.* 2016;253: 66-77. <https://doi.org/10.1016/j.cbi.2016.05.015>.
2. Adil M, Kandhare AD, Visnagri A, Bodhankar SL. Naringin ameliorates sodium arsenite-induced renal and hepatic toxicity in rats: decisive role of KIM-1, Caspase-3, TGF- β , and TNF- α . *Ren Fail.* 2015;37 (8):1396-1407. <https://doi.org/10.3109/0886022X.2015.1074462>
3. Kim YJ, Kim JM. Arsenic Toxicity in Male Reproduction and Development. *Dev Reprod.* 2015;19 (4):167-180. <https://doi.org/10.12717/DR.2015.19.4.167>.
4. Rahmani S, Naraki K, Roohbakhsh A, Hayes AW, Karimi G. The protective effects of rutin on the liver, kidneys, and heart by counteracting organ toxicity caused by synthetic and natural compounds. *Food Sci Nutr.* 2022;11 (1):39-56. <https://doi.org/10.1002/fsn3.3041>.
5. World Health Organization. Obesity and overweight [Internet]. Geneva: World Health Organization; 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
6. Hong YS, Song KH, Chung JY. Health effects of chronic arsenic exposure. *J Prev Med Public Health.* 2014;47(5):245-252. <https://doi.org/10.3961/jpmph.14.035>.
7. Mukherjee AA, Kandhare AD, Bodhankar SL. Elucidation of protective efficacy of Pentahydroxy flavone isolated from *Madhuca indica* against arsenite-induced cardiomyopathy: Role of Nrf-2, PPAR-gamma, c-fos and c-jun. *Environ Toxicol Pharmacol.* 2017;56:172-185. <https://doi.org/10.1016/j.etcp.2017.08.027>
8. Souza AC, Marchesi SC, Ferraz RP, Lima GD, de Oliveira JA, Machado-Neves M. Effects of sodium arsenite and arsenite on male reproductive functions in Wistar rats. *J Toxicol Environ Health A.* 2016;79(6):274-286. <https://doi.org/10.1080/15287394.2016.1150926>.
9. Zubair M, Ahmad M, Qureshi ZI. Review on arsenic-induced toxicity in male reproductive system and its amelioration. *Andrologia.* 2017; 49(9):e12791. <https://doi.org/10.1111/and.12791>.
10. Rachamalla M, Chinthada J, Kushwaha S, Putnala SK, Sahu C, Jena G, et al. Contemporary Comprehensive Review on Arsenic-Induced Male Reproductive Toxicity and Mechanisms of Phytonutrient Intervention. *Toxics.* 2022; 10(12):744. <https://doi.org/10.3390/toxics10120744>.
11. Souza ACE, Bastos DSS, Sertorio MN, Santos FC, Ervilha LOG, de Oliveira LL, et al. Combined effects of arsenic exposure and diabetes on male reproductive functions. *Andrology.* 2019; 7(5):730-740. <https://doi.org/10.1111/andr.12613>.
12. Wang S, Zhao Y, Wu L, Tang M, Su C, Hei TK, Yu Z. Induction of germline cell cycle arrest and apoptosis by sodium arsenite in *Caenorhabditis elegans*. *Chem Res Toxicol.* 2007;20(2):181-186. <https://doi.org/10.1021/tx0601962>.
13. Silva CS, Kudlyk T, Tryndyak VP, Twaddle NC, Robinson B, Gu Q, et al. Gene expression analyses reveal potential mechanism of inorganic arsenic-induced apoptosis in zebrafish. *J Appl Toxicol.* 2023;43(12):1872-1882. <https://doi.org/10.1002/jat.4520>.
14. Uygar R, Aktas C, Caglar V, Uygur E, Erdogan H, Ozen OA. Protective effects of melatonin against arsenic-induced apoptosis and oxidative stress in rat testes. *Toxicol Ind Health.* 2016;32(5):848-859. <https://doi.org/10.1177/0748233713512891>.
15. Omar SM, Zahran NN, Alhotan RA, Hussein EO, Galik B, Saleh AA. Evaluation of *Salvia hispanica* as a Therapeutic Agent against Sodium Arsenic-Induced Testicular Toxicity in a Male Rats Model. *Life (Basel Switzerland).* 2024; 14(1):109. <https://doi.org/10.3390/life14010109>.
16. Pires Das Neves RN, Carvalho F, Carvalho M, Fernandes E, Soares E, de Bastos ML, et al. Protective activity of hesperidin and lipoic acid against sodium arsenite acute toxicity in mice. *Toxicol Pathol.* 2004;32(5):527-535. <https://doi.org/10.1080/01926230490502566>.

17. Negahdari R, Bohlouli S, Sharifi S, Maleki Dizaj S, Rahbar Saadat Y, Khezri K, et al. Therapeutic benefits of rutin and its nanoformulations. *Phytother Res.* 2021; 35(4):1719-1738. <https://doi.org/10.1002/ptr.6904>.
18. Budzynska B, Faggio C, Kruk-Slomka M, Samec D, Nabavi SF, Sureda A, et al. Rutin as Neuroprotective Agent: From Bench to Bedside. *Curr Med Chem.* 2019;26 (27):5152-5164. <https://doi.org/10.2174/0929867324666171003114154>.
19. Hosseinzadeh H, Nassiri-Asl M. Review of the protective effects of rutin on the metabolic function as an important dietary flavonoid. *J Endocrinol Invest.* 2014;37 (9):783-788. <https://doi.org/10.1007/s40618-014-0096-3>.
20. Ganeshpurkar A, Saluja AK. The Pharmaceutical Potential of Rutin. *Saudi Pharm J.* 2017;25 (2):149-164. <https://doi.org/10.1016/j.jps.2016.04.025>
21. Niture NT, Ansari AA, Naik SR. Anti-hyperglycemic activity of rutin in streptozotocin-induced diabetic rats: an effect mediated through cytokines, antioxidants and lipid biomarkers. *Indian J Exp Biol.* 2014;52 (7):720-727. PMID: 25059040.
22. Kamalakkannan N, Prince PS. Antihyperglycaemic and antioxidant effect of rutin, a polyphenolic flavonoid, in streptozotocin-induced diabetic wistar rats. *Basic Clin Pharmacol Toxicol.* 2006;98:97-103. https://doi.org/10.1111/j.1742-7843.2006.pto_241.x
23. Das J, Ghosh J, Manna P, Sinha M, Sil PC. Taurine protects rat testes against NaAsO₂ induced oxidative stress and apoptosis via mitochondrial dependent and independent pathways. *Toxicol Lett.* 2009;187(3):201-210. <https://doi.org/10.1016/j.toxlet.2009.03.001>.
24. Adil M, Visnagri A, Kumar VS, Kandhare AD, Ghosh P, Bodhankar SL. Protective effect of naringin on sodium arsenite induced testicular toxicity via modulation of biochemical perturbations in experimental rats. *Pharmacologia.* 2014;5 (6):222-234. <https://doi.org/10.5567/pharmacologia.2014.222.234>
25. Thangapandiyan S, Miltonprabu S. Epigallocatechin gallate exacerbates fluoride-induced oxidative stress mediated testicular toxicity in rats through the activation of Nrf2 signaling pathway. *Asian Pac J Reprod.* 2015;4 (4):272-287. <https://doi.org/https://doi.org/10.1016/j.apjr.2015.07.005>
26. Kandhare AD, Kumar VS, Adil M, Rajmane AR, Ghosh P, Bodhankar SL. Investigation of gastro protective activity of Xanthium strumarium L. by modulation of cellular and biochemical marker. *Orient Pharm Exp Med.* 2012;12: 287-299. <https://doi.org/10.1007/s13596-012-0090-2>
27. Kandhare AD, Raygude KS, Ghosh P, Ghule AE, Bodhankar SL. Neuroprotective effect of naringin by modulation of endogenous biomarkers in streptozotocin induced painful diabetic neuropathy. *Fitoterapia.* 2012;83 (4):650-659. <https://doi.org/10.1016/j.fitote.2012.01.010>
28. Kandhare AD, Raygude KS, Ghosh P, Ghule AE, Bodhankar SL. Therapeutic role of curcumin in prevention of biochemical and behavioral aberration induced by alcoholic neuropathy in laboratory animals. *Neurosci Lett.* 2012;511(1):18-22. <https://doi.org/10.1016/j.neulet.2012.01.019>
29. Visnagri A, Kandhare AD, Shiva Kumar V, Rajmane AR, Mohammad A, Ghosh P et al. Elucidation of ameliorative effect of Co-enzyme Q10 in streptozotocin-induced diabetic neuropathic perturbation by modulation of electrophysiological, biochemical and behavioral markers. *Biomed Aging Pathol.* 2012;2 (4):157-172. <https://doi.org/http://dx.doi.org/10.1016/j.biomag.2012.10.006>
30. Adil M, Kandhare AD, Dalvi G, Ghosh P, Venkata S, Raygude KS, Bodhankar SL. Ameliorative effect of berberine against gentamicin-induced nephrotoxicity in rats via attenuation of oxidative stress, inflammation, apoptosis and mitochondrial dysfunction. *Ren Fail.* 2016;38 (6):996-1006. <https://doi.org/10.3109/0886022X.2016.1165120>

31. Ketkar S, Rathore A, Kandhare AD, Lohidasan S, Bodhankar SL, Paradkar A, et al. Alleviating exercise-induced muscular stress using neat and processed bee pollen: oxidative markers, mitochondrial enzymes, and myostatin expression in rats. *Integr Med Res.* 2015;4 (3):147-160. <https://doi.org/10.1016/j.imr.2015.02.003>

32. Liu J, Han XC, Wang D, Kandhare AD, Mukherjee-Kandhare AA, Bodhankar SL et al. Elucidation of Molecular Mechanism Involved in Nephroprotective Potential of Naringin in Ethylene Glycol-Induced Urolithiasis in Experimental Uninephrectomized Hypertensive Rats. *Lat Am J Pharm.* 2020;39 (5):991-999.

33. Yao X, Li L, Kandhare AD, Mukherjee-Kandhare AA, Bodhankar SL. Attenuation of reserpine-induced fibromyalgia via ROS and serotonergic pathway modulation by fisetin, a plant flavonoid polyphenol. *Exp Ther Med.* 2020;19 (2):1343-1355. <https://doi.org/10.3892/etm.2019.8328>

34. Zhang K, Kandhare A, Mukherjee-Kandhare A, Bodhankar S. Apigenin attenuated ethylene glycol induced urolithiasis in uninephrectomized hypertensive rats: A possible role of bikunin, BMP-2/4, and osteopontin. *Pharmacogn Mag.* 2020;16 (71):455-463. https://doi.org/10.4103/pm.pm_83_20

35. McNamara ER, Madden-Fuentes RJ, Routh JC, Rouse D, Madden JF, Wiener JS, et al. Evaluation of cold ischemia for preservation of testicular function during partial orchiectomy in the rat model. *J Pediatr Urol.* 2014;10 (4):593-597. <https://doi.org/10.1016/j.jpurol.2014.06.005>

36. Prathima P, Pavani R, Sukeerthi S, Sainath SB. α -Lipoic acid inhibits testicular and epididymal oxidative damage and improves fertility efficacy in arsenic-intoxicated rats. *J Biochem Mol Toxicol.* 2018;32 (2):e22016. <https://doi.org/10.1002/jbt.22016>.

37. Huang Q, Luo L, Alamdar A, Zhang J, Liu L, Tian M, et al. Integrated proteomics and metabolomics analysis of rat testis: Mechanism of arsenic-induced male repro-ductive toxicity. *Sci Rep.* 2016;6:32518. <https://doi.org/10.1038/srep32518>.

38. Pant N, Murthy RC, Srivastava SP. Male reproductive toxicity of sodium arsenite in mice. *Hum Exp Toxicol.* 2004; 23(8):399-403. <https://doi.org/10.1191/0960327104ht467oa>.

39. Oduwole OO, Huhtaniemi IT, Misrahi M. The Roles of Luteinizing Hormone, Follicle-Stimulating Hormone and Testosterone in Spermatogenesis and Folliculogenesis Revisited. *Int J Mol Sci.* 2021;22 (23):12735. <https://doi.org/10.3390/ijms222312735>

40. Oduwole OO, Peltoketo H, Huhtaniemi IT. Role of Follicle-Stimulating Hormone in Spermatogenesis. *Front Endocrinol. (Lausanne).* 2018;9:763. <https://doi.org/10.3389/fendo.2018.00763>

41. Prakash C, Soni M, Kumar V. Mitochondrial oxidative stress and dysfunction in arsenic neurotoxicity: A review. *J Appl Toxicol.* 2016; 36 (2):179-188. <https://doi.org/10.1002/jat.3256>

42. Keshavarz-Bahaghighe H, Sepand MR, Ghahremani MH, Aghsami M, Sanadgol N, Omidi A, et al. Acetyl-L-Carnitine Attenuates Arsenic-Induced Oxidative Stress and Hippocampal Mitochondrial Dysfunction. *Biol Trace Elem Res.* 2018;184 (2):422-435. <https://doi.org/10.1007/s12011-017-1210-0>

43. Thakur M, Rachamalla M, Niyogi S, Datusalia AK, Flora SJS. Molecular Mechanism of Arsenic-Induced Neurotoxicity including Neuronal Dysfunctions. *Int J Mol Sci.* 2021;22 (18):10077. <https://doi.org/10.3390/ijms221810077>

44. Liu L, Trimarchi JR, Navarro P, Blasco MA, Keefe DL. Oxidative Stress Contributes to Arsenic-induced Telomere Attrition, Chromosome Instability, and Apoptosis. *J Biol Chem.* 2003;278 (34):31998-32004. <https://doi.org/10.1074/jbc.m303553200>

45. Cui J, Wang G, Kandhare AD, Mukherjee-Kandhare AA, Bodhankar SL. Neuroprotective effect of naringin, a flavone glycoside in quinolinic acid-induced neurotoxicity: Possible role of PPAR-gamma, Bax/Bcl-2, and caspase-3. *Food Chem.*

Toxicol. 2018;121:95-108. <https://doi.org/10.1016/j.jct.2018.08.028>

46. Kandhare AD, Mukherjee AA, Bodhankar SL. Neuroprotective effect of *Azadirachta indica* standardized extract in partial sciatic nerve injury in rats: Evidence from anti-inflammatory, antioxidant and anti-apoptotic studies. EXCLI J. 2017;16:546-565. <https://doi.org/doi: 10.17179/excli2017-161>.

47. Raygude KS, Kandhare AD, Ghosh P, Ghule AE, Bodhankar SL. Evaluation of ameliorative effect of quercetin in experimental model of alcoholic neuropathy in rats. Inflammopharmacol. 2012; 20 (6):331-341. <https://doi.org/10.1007/s10787-012-0122-z>

48. Xu M, Rui D, Yan Y, Xu S, Niu Q, Feng G, et al. Oxidative Damage Induced by Arsenic in Mice or Rats: A Systematic Review and Meta-Analysis. Biol Trace Elem Res. 2017; 76 (1):154-175. <https://doi.org/10.1007/s12011-016-0810-4>

49. Visnagri A, Kandhare AD, Chakravarty S, Ghosh P, Bodhankar SL. Hesperidin, a flavanoglycone attenuates experimental diabetic neuropathy via modulation of cellular and biochemical marker to improve nerve functions. Pharm Biol. 2014;52 (7):814-828. <https://doi.org/10.3109/13880209.2013.870584>

50. Jomova K, Jenisova Z, Feszterova M, Barros S, Liska J, Hudecova D, et al. Arsenic: toxicity, oxidative stress and human disease. J Appl Toxicol. 2011;31 (2):95-107. <https://doi.org/10.1002/jat.1649>

51. Biswas J, Sinha D, Mukherjee S, Roy S, Siddiqi M, Roy M. Curcumin protects DNA damage in a chronically arsenic-exposed population of West Bengal. Hum Exp Toxicol. 2010;29 (6):513-524. <https://doi.org/10.1177/0960327109359020>

52. Almeer R, Alyami NM. The protective effect of apigenin against inorganic arsenic salt-induced toxicity in PC12 cells. Environ Sci Poll Res. 2023;30:106625-106635. <https://doi.org/10.1007/s11356-023-29884-w>

53. El-Khadragy M, Al-Meghrin WA, Alsadhan NA, Metwally DM, El-Hennamy RE, Sa-lem FEH, et al. Impact of Coenzyme Q10 Administration on Lead Acetate-Induced Testicular Damage in Rats. Oxid Med Cell Longev. 2020;2020: 4981386. <https://doi.org/10.1155/2020/4981386>

54. Devkar ST, Kandhare AD, Zanwar AA, Jagtap SD, Katyare SS, Bodhankar SL, et al. Hepatoprotective effect of withanolide-rich fraction in acetaminophen-intoxicated rat: decisive role of TNF- α , IL-1 β , COX-II and iNOS. Pharm Biol. 2016; 54 (11): 2394-2403. <https://doi.org/10.3109/13880209.2016.1157193>.

55. Honmore VS, Kandhare AD, Kadam PP, Khedkar VM, Natu AD, Rojatkar SR, et al. Diarylheptanoid, a constituent isolated from methanol extract of *Alpinia officinarum* attenuates TNF-alpha level in Freund's complete adjuvant-induced arthritis in rats. J Ethnopharmacol. 2019;229:233-245. <https://doi.org/10.1016/j.jep.2018.10.019>

56. Kandhare AD, Bodhankar SL, Mohan V, Thakurdesai PA. Effect of glycosides based standardized fenugreek seed extract in bleomycin-induced pulmonary fibrosis in rats: Decisive role of Bax, Nrf2, NF- κ B, Muc5ac, TNF- α and IL-1 β . Chem Biol Interact. 2015; 237:151-165. <https://doi.org/10.1016/j.cbi.2015.06.019>.

57. Rao G, Qiao B, Zhong G, Li T, Su Q, Wu S, et al. Arsenic and polystyrene-nano plastics co-exposure induced testicular toxicity: Triggers oxidative stress and promotes apoptosis and inflammation in mice. Environ Toxicol. 2023;39 (1):264-276. <https://doi.org/10.1002/tox.23970>

58. Joza N, Susin SA, Daugas E, Stanford WL, Cho SK, Li CY, et al. Essential role of the mitochondrial apoptosis-inducing factor in programmed cell death. Nature 2001;410(6828):549-554. <https://doi.org/10.1038/35069004>

59. Zhao P, Wang Y, Yang Q, Yu G, Ma F, Dong J. Abamectin causes cardiac dysfunction in carp via inhibiting redox equilibrium and resulting in immune inflammatory response and programmed cell death. Environ Sci Pollut Res Int. 2023;30 (11):29494-29509. <https://doi.org/10.1007/s11356-022-24004-6>

60. **Ma T, Kandhare AD, Mukherjee-Kandhare AA, Bodhankar SL.** Fisetin, a plant flavonoid ameliorates doxorubicin-induced cardiotoxicity in experimental rats: the decisive role of caspase-3, COX-II, cTn-I, iNOs and TNF-alpha. *Mol Biol Rep* 2019;46 (1):105-118. <https://doi.org/10.1007/s11033-018-4450-y>

61. **Visnagri A, Kandhare AD, Bodhankar SL.** Renoprotective effect of berberine via inhibition on apoptosis and mitochondrial-dependent pathway in renal ischemia reperfusion-induced mutilation. *Ren Fail*. 2015;37(3):482-493. <https://doi.org/10.3109/0886022X.2014.996843>

62. **Ghule AE, Kandhare AD, Jadhav SS, Zanwar AA, Bodhankar SL.** Omega-3-fatty acid adds to the protective effect of flax lignan concentrate in pressure overload-induced myocardial hypertrophy in rats via modulation of oxidative stress and apoptosis. *Int Immunopharmacol*. 2015; 28 (1):751-763. <https://doi.org/10.1016/j.intimp.2015.08.005>.